



REC'D 15 AUG 2000

WIPO

PCT

Kongeriget Danmark

DK 00/00413

Patent application No.: PA 1999 01014

Date of filing: 20 July 1999

Applicant: M&E Biotech A/S
Kogle Allé 6
DK-2970 Hørsholm

This is to certify the correctness of the following information:

The attached photocopy is a true copy of the following document:

- The specification, claims and drawings as filed with the application on the filing date indicated above.



PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)



Patent- og
Varemærkestyrelsen
Erhvervsministeriet

Taastrup, 01 August 2000


Karin Schlichting
Head Clerk

20 JUL 1999

FIELD OF THE INVENTION

The present invention generally relates to the field of animal and poultry husbandry. More specifically, the present invention
5 pertains to improvements in controlling and increasing growth in farm animals for slaughter. Thus, provided are novel methods and means for increasing growth in animals for slaughter.

BACKGROUND OF THE INVENTION

Currently, two different medical approaches are used in order to
10 increase the growth rate of farm animals: On the one hand the administration to farm animals of antibiotics or antibiotic-like compounds, and on the other the administration of growth hormones.

The administration of antibiotics and antibiotic-like compounds
15 in farm animals (notably pigs) in order to promote the growth rate of these animals has in later years been proven to cause several problems. Some of these compounds are chemically closely related to antibiotics used for treatment of human disease, and evidence is building up that extensive use of such compounds in
20 farm animals induces cross-resistance towards human antibiotics in microorganisms pathogenic in man. Further, relatively low increases in growth rates of 1-3% are obtained with these compounds.

The use of growth hormones in farm animals is expensive, and the
25 treatment has to be repeated at rather short intervals due to the relatively short half-life of growth hormone. Furthermore, the presence of potential residual growth hormone in meat produced from treated animals has created some concerns in European consumers particularly.

GDF-8

GDF-8 or myostatin was discovered in May 1997 as a growth regulating factor selectively down-regulating skeletal muscle growth (McPherron et al., Nature, 387, 83-90, 1997). GDF-8 expression is restricted to the myotome compartment of developing embryonal somites, but it is also expressed in various muscle tissues throughout the body in the adult animal.

GDF-8 knock-out mice exhibit a strongly increased skeletal muscle mass. The increase in skeletal muscle mass appear to be widespread throughout the body, and isolated muscles from GDF-8 negative mice weigh about 2-3 times more than wild type muscles. The total body weights of the knock-out mice are about 35% higher than wild type mice and mice lacking the GDF-8 gene has more than 80% more muscle fibres compared to normal mice. The massive skeletal muscle enlargement observed in the knock-out mice is, however, not merely due to an increase in muscle fibre numbers but also to a significant muscle fibre hypertrophy. The cross-sectional muscle area of the GDF-8 knock-out mice is increased by about 14 to 49% depending on muscle type.

Interestingly, there is also observed an enhanced rate of muscle mass increase in adult transgenic mice compared to adult non-transgenic mice. Further, all GDF-8 negative mice has been shown to be viable and fertile.

In November 1997 the authors who originally discovered GDF-8 published that two breeds of cattle that are characterized by strongly increased muscle mass, Belgian Blue and Piedmontese, have mutations in the GDF-8 coding sequence and that this accounts for their large muscles (McPherron and Se-Jin Lee, 1997, PNAS 94, 12457-12461). This phenomenon of "double muscling" has been observed in many breeds of cattle for the past 190 years, and the animals have an average increase in

muscle mass of 20-25%. They also show an increased feed efficiency, but they still produce high-quality meat.

Unlike the GDF-8 knock-out mice, however, the Belgian Blue cattle also exhibit a reduction in mass of most other organs.

- 5 These "natural knock-out cows" also suffer from reduced female fertility, reduced viability of offspring, and a delayed sexual maturation.
-

The relative increase in muscle mass in "knock-out cows" is not nearly as pronounced as is observed in the knock-out mice - in
10 fact, it corresponds to the extent of muscle hypertrophy observed in the mice. McPherron et al. speculate that one reason could be that normal cattle may be nearer than mice to a maximum limit of muscle size (and hence to the maximum obtainable number of muscle fibres) after generations of selective breeding. No
15 data regarding the number of muscle fibre hyperplasia versus hypertrophy in cattle was published by the authors, but based on this assumption and the muscle hypertrophy observed in the knock-out mice, it could be possible that the increase in muscle mass and growth rate observed in e.g. the Belgian Blue cattle is
20 largely due to muscle hypertrophy and to a lesser extent muscle hyperplasia.

Physiological role of GDF-8

Expression of GDF-8 is highly restricted to skeletal muscle. There is a low level of expression in adipose tissue, but
25 notably there is no expression in heart muscle. The physiological role of GDF-8 in the adult individual is not known, although it seems that GDF-8 may function as a specific negative regulator of skeletal muscle growth. Speculations about the physiological role centres upon important functions in
30 exercise induced muscle hypertrophy or regeneration after muscle injury. GDF-8 may, however, also suppress adipose tissue growth.

It is not known whether GDF-8 works locally or systemically during the growth of the animal.

Structure of GDF-8

GDF-8 belongs to the transforming growth factor β (TGF- β) superfamily which encompasses a group of structurally-related proteins involved in embryonic development. Human and bovine GDF-8 are produced as 375 amino acids long precursor proteins. As other TGF- β superfamily proteins GDF-8 is probably processed proteolytically into a much shorter C-terminal fragment of about 109 amino acids which form disulphide linked homodimers. The homodimer is probably the biologically active form of GDF-8.

The amino acid sequences of murine, rat, human, baboon, bovine, porcine, sheep, chicken and turkey GDF-8 are known and the GDF-8 molecule is highly conserved (McPherron and Se-Jin Lee, PNAS, 94, 12457-12461, 1997). The sequences of murine, rat, human, porcine, chicken and turkey GDF-8 are 100% identical in the C-terminal region, which probably contains the biologically active part of GDF-8. Both bovine and sheep GDF-8 only differ by two amino acid residues from human GDF-8 in the C-terminal region. Bovine GDF-8 has the sequence -Glu-Gly- instead of -Lys-Glu- in positions 356-357, and sheep GDF-8 has two conservative substitutions (a Val in position 316 and a Arg in position 333 instead of Leu and Lys, respectively). None of the known GDF-8 proteins include potential N-glycosylation sites in their active C-terminal region.

OBJECT OF THE INVENTION

It is an object of the present invention to provide a recombinant therapeutic vaccine capable of effecting down-regulation of growth differentiation factor 8 (GDF-8) (also known as myostatin) in order to increase the muscle growth rate

of farm animals. It is a further object to provide a method of vaccinating animals in order to increase the muscle growth rate. It is also an object to provide variants of GDF-8 which are capable of breaking autotolerance against autologous GDF-8. Yet
5 a further object is to provide nucleic acid fragments, vectors and transformed cells which are all useful in the preparation of the vaccines and GDF-8 variants.

SUMMARY OF THE INVENTION

Based on the above-referenced findings in transgenic GDF-8
10 knock-out animals and in Belgian Blue and Piedmontese, the present inventors have developed the theory that it will be possible to down-regulate GDF-8 in animals by inducing an effective and fine-tuned immune response against GDF-8. Hence, what is basically provided by the present invention is a method
15 and means to immunologically down-regulate GDF-8 in order to effect an increase the muscle mass in animals.

The advantages of this approach over known growth-enhancing measures in farm animals are several. First of all, problems relating to cross-resistance in pathological microorganisms will
20 not arise when using the present approach. Further, there will be no potential residual exogenously administered growth hormone in meat from animals which have been subjected to the present treatment. Finally, the ethical problems involved in the breeding and production of cattle like the Belgian Blue and the
25 Piedmontese (where many calves are born by means of Caesarian section and where other organs are of reduced size) can be completely avoided by postponing the down-regulation of GDF-8 in the animals until after birth (e.g. in adult life only) - in fact, animals having an increased muscle mass need not give
30 birth at all, and the treatment can therefore be reserved to those animals which are predestined to become slaughtered.

Since the number and type of muscle fibres is determined during embryonic life, it is not very likely that an anti-GDF-8 vaccine would increase the number of muscle fibres in adult farm animals. It is therefore neither certain nor likely that an anti-GDF-8 vaccine would be able to suppress GDF-8 to the same level as seen in GDF-8 knock-out animals. This would probably not be desirably either, since this perhaps could negatively affect meat quality, breeding capability, and fat ratio.

Nevertheless, an increase in growth rate and/or maximum body weight of 5-25 % obtained in vaccinated animals due to post partum muscle hypertrophy (a figure range which does not seem unrealistic) would still be interesting in the production of meat from cows, pigs and poultry.

The sequence identity of GDF-8 relative to other members of the TGF- β family is only 30-40 percent at the amino acid level. This low sequence identity will in all likelihood not create problems with cross-reactivity of antibodies induced towards GDF-8.

Thus, in its broadest and most general scope, the present invention relates to a method for *in vivo* down-regulation of growth differentiation factor 8 (GDF-8) activity in an animal, including a human being, the method comprising effecting presentation to the animal's immune system of an immunologically effective amount of

- at least one GDF-8 polypeptide or subsequence thereof which has been formulated so that immunization of the animal with the GDF-8 polypeptide or subsequence thereof induces production of antibodies against the GDF-8 polypeptide, and/or
- at least one GDF-8 analogue wherein is introduced at least one modification in the GDF-8 amino acid sequence which has as a result that immunization of the animal with the analogue induces production of antibodies against the GDF-8 polypeptide.

It is expected that 1-4 annual injections with an immunogenic composition according to the invention will be sufficient to obtain the desired effect, whereas administration of both growth hormones and antibiotics require much more frequent
 5 administrations to the animal in question.

The invention also relates to GDF-8 analogues as well as to nucleic acid fragments encoding a subset of these. Also

 immunogenic compositions comprising the analogues or the nucleic acid fragments are part of the invention.

10 The invention also relates to a method of identifying immunogenically effective analogues of GDF-8 as well as a method for preparing composition comprising the GDF-8 analogues.

LEGENDS TO THE FIGURES

Fig. 1: Models of GDF-8 derived autovaccine constructs.

15 Stretches indicated in dark grey are the stretches in which substitutions with P2 and P30, respectively, are proposed to take place.

A: Monomer construct with P2 epitope inserts. "1" indicates a P2 substitution of amino acid residues 18-32 of the 109 aa C-
 20 terminal GDF-8 fragment, "2" indicates a P2 substitution of amino acid residues 52-66 of the 109 aa C-terminal GDF-8 fragment, and "3" indicates a P2 substitution of amino acid residues 83-97 of the 109 aa C-terminal GDF-8 fragment.

B: Monomer construct with P30 epitope inserts. "1" indicates a
 25 P30 substitution of amino acid residues 21-41 of the 109 aa C-terminal GDF-8 fragment, "2" indicates a P30 substitution of amino acid residues 49-69 of the 109 aa C-terminal GDF-8 fragment, and "3" indicates a P30 substitution of amino acid residues 79-99 or 84-104 of the 109 aa C-terminal GDF-8
 30 fragment.

C: Extended monomer construct with P2 and P30 substituted into the 160 aa C-terminal GDF-8 fragment.

D: Dimer construct with the P2 and P30 epitopes serially linked between two copies of the 109 C-terminal GDF-8 fragment.

5 Fig. 2: Expected 3D structure of the TGF- β protein.

The model is based on an analogous domain (TGJ-1) from the TGF- β protein. β -sheets are shown as arrows and helices as cylinders.

DETAILED DISCLOSURE OF THE INVENTION

Definitions

10 In the following, a number of terms used in the present specification and claims will be defined and explained in detail in order to clarify the metes and bounds of the invention.

The terms "T-lymphocyte" and "T-cell" will be used interchangeably for lymphocytes of thymic origin which are
15 responsible for various cell mediated immune responses as well as for helper activity in the humoral immune response. Likewise, the terms "B-lymphocyte" and "B-cell" will be used interchangeably for antibody-producing lymphocytes.

A "GDF-8 polypeptide" is herein intended to denote polypeptides
20 having the amino acid sequence of the above-discussed GDF-8 proteins derived from a number of animals (or truncates thereof sharing a substantial amount of B-cell epitopes with intact GDF-8), but also polypeptides having the amino acid sequence identical to xeno-analogues of these two proteins isolated from
25 other species are embraced by the term. When using the term is normally meant the biologically active form, i.e. the C-terminal peptide which in humans is of 109 amino acids in length. Also unglycosylated forms of GDF-8 which are prepared in prokaryotic system are included within the boundaries of the term as are

forms having varying O-glycosylation patterns due to the use of e.g. yeasts or other non-mammalian eukaryotic expression systems. It should, however, be noted that when using the term "a GDF-8 polypeptide" it is intended that the polypeptide in question is normally non-immunogenic when presented to the animal to be treated. In other words, the GDF-8 polypeptide is a self-protein or is a xeno-analogue of such a self-protein which will not normally give rise to an immune response against GDF-8 of the animal in question.

10 A "GDF-8 analogue" is a GDF-8 polypeptide which has been subjected to changes in its primary structure. Such a change can e.g. be in the form of fusion of a GDF-8 polypeptide to a suitable fusion partner (i.e. a change in primary structure exclusively involving C- and/or N-terminal additions of amino acid residues) and/or it can be in the form of insertions and/or deletions and/or substitutions in the GDF-8 polypeptide's amino acid sequence. Also derivatized GDF-8 molecules are encompassed by the term a "GDF-8 analogue", cf. the discussion below which deals with modifications of GDF-8.

20 It should be noted that the use as a vaccine in a human of e.g. a canine analogue of human GDF-8 can be imagined to produce the desired immunity against GDF-8. Such use of an xeno-analogue for immunization is also considered to be an "GDF-8 analogue" as defined above.

25 The term "polypeptide" is in the present context intended to mean both short peptides of from 2 to 10 amino acid residues, oligopeptides of from 11 to 100 amino acid residues, and polypeptides of more than 100 amino acid residues. Furthermore, the term is also intended to include proteins, i.e. functional biomolecules comprising at least one polypeptide; when comprising at least two polypeptides, these may form complexes, be covalently linked, or may be non-covalently linked. The

polypeptide(s) in a protein can be glycosylated and/or lipidated and/or comprise prosthetic groups.

The term "subsequence" means any consecutive stretch of at least 3 amino acids or, when relevant, of at least 3 nucleotides,
5 derived directly from a naturally occurring GDF-8 amino acid sequence or nucleic acid sequence, respectively.

The term "animal" is in the present context in general intended to denote an animal species (preferably mammalian), such as *Homo sapiens*, *Canis domesticus*, etc. and not just one single animal.
10 However, the term also denotes a population of such an animal species, since it is important that the individuals immunized according to the method of the invention all harbour substantially the same GDF-8 allowing for immunization of the animals with the same immunogen(s). If, for instance, genetic
15 variants of GDF-8 exist in a different population it may be necessary to use different immunogens in these different populations in order to be able to break the autotolerance towards GDF-8 in each population. It will be clear to the skilled person that an animal in the present context is a living
20 being which has an immune system. It is preferred that the animal is a vertebrate, such as a mammal.

By the term "in vivo down-regulation of GDF-8 activity" is herein meant reduction in the living organism of the number of interactions between GDF-8 and its receptors (or between GDF-8
25 and other possible biologically important binding partners for this molecule). The down-regulation can be obtained by means of several mechanisms: Of these, simple interference with the active site in GDF-8 by antibody binding is the most simple. However, it is also within the scope of the present invention
30 that the antibody binding results in removal of GDF-8 by scavenger cells (such as macrophages and other phagocytic cells).

The expression "effecting presentation ... to the immune system" is intended to denote that the animal's immune system is subjected to an immunogenic challenge in a controlled manner. As will appear from the disclosure below, such challenge of the immune system can be effected in a number of ways of which the most important are vaccination with polypeptide containing "pharmaccines" (i.e. a vaccine which is administered to treat or ameliorate ongoing disease) or nucleic acid "pharmaccine"

vaccination. The important result to achieve is that immune competent cells in the animal are confronted with the antigen in an immunologically effective manner, whereas the precise mode of achieving this result is of less importance to the inventive idea underlying the present invention.

The term "immunogenically effective amount" has its usual meaning in the art, i.e. an amount of an immunogen which is capable of inducing an immune response which significantly engages pathogenic agents which share immunological features with the immunogen.

When using the expression that the GDF-8 has been "modified" is herein meant a chemical modification of the polypeptide which constitutes the backbone of GDF-8. Such a modification can e.g. be derivatization (e.g. alkylation, acylation, esterification etc.) of certain amino acid residues in the GDF-8 sequence, but as will be appreciated from the disclosure below, the preferred modifications comprise changes of (or additions to) the primary structure of the GDF-8 amino acid sequence, i.e. modifications which results in the provision of a GDF-8 analogue where the primary amino acid sequence has been modified.

When discussing "autotolerance towards GDF-8" is understood that since GDF-8 is a self-protein in the population to be vaccinated, normal individuals in the population do not mount an immune response against GDF-8; it cannot be excluded, though, that occasional individuals in an animal population might be

able to produce antibodies against native GDF-8, e.g. as part of an autoimmune disorder. At any rate, an animal will normally only be autotolerant towards its own GDF-8, but it cannot be excluded that GDF-8 analogues derived from other animal species or from a population having a different GDF-8 phenotype would also be tolerated by said animal.

A "foreign T-cell epitope" (or: "foreign T-lymphocyte epitope") is a peptide which is able to bind to an MHC molecule and which stimulates T-cells in an animal species. Preferred foreign T-cell epitopes in the invention are "promiscuous" epitopes, i.e. epitopes which bind to a substantial fraction of a particular class of MHC molecules in an animal species or population. Only a very limited number of such promiscuous T-cell epitopes are known, and they will be discussed in detail below. It should be noted that in order for the immunogens which are used according to the present invention to be effective in as large a fraction of an animal population as possible, it may be necessary to 1) insert several foreign T-cell epitopes in the same GDF-8 analogue or 2) prepare several GDF-8 analogues wherein each analogue has a different promiscuous epitope inserted. It should be noted also that the concept of foreign T-cell epitopes also encompasses use of cryptic T-cell epitopes, i.e. epitopes which are derived from a self-protein and which only exerts immunogenic behaviour when existing in isolated form without being part of the self-protein in question.

A "foreign T helper lymphocyte epitope" (a foreign T_H epitope) is a foreign T cell epitope which binds an MHC Class II molecule and can be presented on the surface of an antigen presenting cell (APC) bound to the MHC Class II molecule.

A "functional part" of a (bio)molecule is in the present context intended to mean the part of the molecule which is responsible for at least one of the biochemical or physiological effects exerted by the molecule. It is well-known in the art that many

enzymes and other effector molecules have an active site which is responsible for the effects exerted by the molecule in question. Other parts of the molecule may serve a stabilizing or solubility enhancing purpose and can therefore be left out if
5 these purposes are not of relevance in the context of a certain embodiment of the present invention. For instance it is possible to use certain other cytokines as a modifying moiety in GDF-8 (cf. the detailed discussion below), and in such a case, the

issue of stability may be irrelevant since the coupling to GDF-8
10 provides the stability necessary.

The term "adjuvant" has its usual meaning in the art of vaccine technology, *i.e.* a substance or a composition of matter which is
1) not in itself capable of mounting an immune response against the immunogen of the vaccine, but which is 2) nevertheless
15 capable of enhancing the immune response against the immunogen. Or, in other words, vaccination with the adjuvant alone does not provide an immune response against the immunogen, vaccination with the immunogen may or may not give rise to an immune response against the immunogen, but the combined vaccination
20 with immunogen and adjuvant induces an immune response against the immunogen which is furthermore stronger than that induced by the immunogen alone.

"Targeting" of a molecule is in the present context intended to denote the situation where a molecule upon introduction in the
25 animal will appear preferentially in certain tissue(s) or will be preferentially associated with certain cells or cell types. The effect can be accomplished in a number of ways including formulation of the molecule in composition facilitating targeting or by introduction in the molecule of groups which
30 facilitates targeting. These issues will be discussed in detail below.

"Stimulation of the immune system" means that a substance or composition of matter exhibits a general, non-specific

immunostimulatory effect. A number of adjuvants and putative adjuvants (such as certain cytokines) share the ability to stimulate the immune system. The result of using an immunostimulating agent is an increased "alertness" of the immune system meaning that simultaneous or subsequent immunization with an immunogen induces a significantly more effective immune response compared to isolated use of the immunogen.

Preferred embodiments of GDF-8 activity down-regulation

- 10 It is preferred that the GDF-8 polypeptide used as an immunogen in the method of the invention is a modified molecule wherein at least one change is present in the GDF-8 amino acid sequence, since the chances of obtaining the all-important breaking of autotolerance towards GDF-8 is greatly facilitated that way. It should be noted that this does not exclude the possibility of using such a modified GDF-8 in formulations which further facilitate the breaking of autotolerance against GDF-8, e.g. formulations containing certain adjuvants discussed in detail below.
- 20 It has been shown (in Dalum I et al., 1996, J. Immunol. 157: 4796-4804) that potentially self-reactive B-lymphocytes recognizing self-proteins are physiologically present in normal individuals. However, in order for these B-lymphocytes to be induced to actually produce antibodies reactive with the relevant self-proteins, assistance is needed from cytokine producing T-helper lymphocytes (T_H -cells or T_H -lymphocytes). Normally this help is not provided because T-lymphocytes in general do not recognize T-cell epitopes derived from self-proteins when presented by antigen presenting cells (APCs).
- 30 However, by providing an element of "foreignness" in a self-protein (i.e. by introducing an immunologically significant modification), T-cells recognizing the foreign element are activated upon recognizing the foreign epitope on an APC (such

as, initially, a mononuclear cell). Polyclonal B-lymphocytes (which are also specialised APCs) capable of recognising self-epitopes on the modified self-protein also internalise the antigen and subsequently presents the foreign T-cell epitope(s) thereof, and the activated T-lymphocytes subsequently provide cytokine help to these self-reactive polyclonal B-lymphocytes. Since the antibodies produced by these polyclonal B-lymphocytes are reactive with different epitopes on the modified polypeptide, including those which are also present in the native polypeptide, an antibody cross-reactive with the non-modified self-protein is induced. In conclusion, the T-lymphocytes can be led to act as if the population of polyclonal B-lymphocytes have recognised an entirely foreign antigen, whereas in fact only the inserted epitope(s) is/are foreign to the host. In this way, antibodies capable of cross-reacting with non-modified self-antigens are induced.

Several ways of modifying a peptide self-antigen in order to obtain breaking of autotolerance are known in the art. Hence, according to the invention, the modification can include that

- 20 - at least one foreign T-cell epitope is introduced, and/or
- at least one first moiety is introduced which effects targeting of the modified molecule to an antigen presenting cell (APC), and/or
- at least one second moiety is introduced which stimulates the immune system, and/or
- 25 - at least one third moiety is introduced which optimizes presentation of the modified GDF-8 polypeptide to the immune system.

However, all these modifications should be carried out while maintaining a substantial fraction of the original B-lymphocyte epitopes in active GDF-8, since the B-lymphocyte recognition of the native molecule is thereby enhanced.

In one preferred embodiment, side groups (in the form of foreign T-cell epitopes or the above-mentioned first, second and third moieties) are covalently or non-covalently introduced. This is intended to mean that stretches of amino acid residues derived from GDF-8 are derivatized without altering the primary amino acid sequence, or at least without introducing changes in the peptide bonds between the individual amino acids in the chain.

An alternative, and preferred, embodiment utilises amino acid substitution and/or deletion and/or insertion and/or addition (which may be effected by recombinant means or by means of peptide synthesis; modifications which involves longer stretches of amino acids can give rise to fusion polypeptides). One especially preferred version of this embodiment is the technique described in WO 95/05849, which discloses a method for down-regulating self-proteins by immunising with analogues of the self-proteins wherein a number of amino acid sequence(s) has been substituted with a corresponding number of amino acid sequence(s) which each comprise a foreign immunodominant T-cell epitope, while at the same time maintaining the overall tertiary structure of the self-protein in the analogue. For the purposes of the present invention, it is however sufficient if the modification (be it an amino acid insertion, addition, deletion or substitution) gives rise to a foreign T-cell epitope and at the same time preserves a substantial number of the B-cell epitopes in GDF-8. However, in order to obtain maximum efficacy of the immune response induced, it is preferred that the overall tertiary structure of GDF-8 is maintained in the modified molecule.

The following formula describes the GDF-8 constructs generally covered by the invention:

$$(\text{MOD}_1)_{s1} (\text{GDF-8}_{e1})_{n1} (\text{MOD}_2)_{s2} (\text{GDF-8}_{e2})_{n2} \dots (\text{MOD}_x)_{sx} (\text{GDF-8}_{ex})_{nx} \quad (\text{I})$$

-where GDF-8_{e1}-GDF-8_{ex} are x B-cell epitope containing subsequences of GDF-8 which independently are identical or non-identical and which may contain or not contain foreign side groups, x is an integer ≥ 3 , n1-nx are x integers ≥ 0 (at least one is ≥ 1), MOD₁-MOD_x are x modifications introduced between the preserved B-cell epitopes, and s₁-s_x are x integers ≥ 0 (at least one is ≥ 1 if no side groups are introduced in the GDF-8_e sequences). Thus, given the general functional restraints on the immunogenicity of the constructs, the invention allows for all kinds of permutations of the original GDF-8 sequence, and all kinds of modifications therein. Thus, included in the invention are modified GDF-8 obtained by omission of parts of the GDF-8 sequence which e.g. exhibit adverse effects *in vivo* or omission of parts which could give rise to undesired immunological reactions.

Maintenance of a substantial fraction of B-cell epitopes or even the overall tertiary structure of a protein which is subjected to modification as described herein can be achieved in several ways. One is simply to prepare a polyclonal antiserum directed against GDF-8 (e.g. an antiserum prepared in a rabbit) and thereafter use this antiserum as a test reagent (e.g. in a competitive ELISA) against the modified proteins which are produced. Modified versions (analogues) which react to the same extent with the antiserum as does GDF-8 must be regarded as having the same overall tertiary structure as GDF-8 whereas analogues exhibiting a limited (but still significant and specific) reactivity with such an antiserum are regarded as having maintained a substantial fraction of the original B-cell epitopes.

Alternatively, a selection of monoclonal antibodies reactive with distinct epitopes on GDF-8 can be prepared and used as a test panel. This approach has the advantage of allowing 1) an epitope mapping of GDF-8 and 2) a mapping of the epitopes which are maintained in the analogues prepared.

Of course, a third approach would be to resolve the 3-dimensional structure of GDF-8 or of a biologically active truncate thereof (cf. above) and compare this to the resolved three-dimensional structure of the analogues prepared. Three-dimensional structure can be resolved by the aid of X-ray diffraction studies and NMR-spectroscopy. Further information relating to the tertiary structure can to some extent be obtained from circular dichroism studies which have the advantage of merely requiring the polypeptide in pure form (whereas X-ray diffraction requires the provision of crystallized polypeptide and NMR requires the provision of isotopic variants of the polypeptide) in order to provide useful information about the tertiary structure of a given molecule. However, ultimately X-ray diffraction and/or NMR are necessary to obtain conclusive data since circular dichroism can only provide indirect evidence of correct 3-dimensional structure via information of secondary structure elements.

One preferred embodiment of the invention utilises multiple presentations of B-lymphocyte epitopes of GDF-8 (i.e. formula I wherein at least one B-cell epitope is present in two positions). This effect can be achieved in various ways, e.g. by simply preparing fusion polypeptides comprising the structure $(\text{GDF-8})_m$, where m is an integer ≥ 2 and then introduce the modifications discussed herein in at least one of the GDF-8 sequences, or alternatively, inserted between at least two of the GDF-8 amino acid sequences. It is preferred that the modifications introduced includes at least one duplication of a B-lymphocyte epitope and/or the introduction of a hapten.

As mentioned above, the introduction of a foreign T-cell epitope can be accomplished by introduction of at least one amino acid insertion, addition, deletion, or substitution. Of course, the normal situation will be the introduction of more than one change in the amino acid sequence (e.g. insertion of or substitution by a complete T-cell epitope) but the important goal

to reach is that the GDF-8 analogue, when processed by an antigen presenting cell (APC), will give rise to such a foreign immunodominant T-cell epitope being presented in context of an MCH Class II molecule on the surface of the APC. Thus, if the
5 GDF-8 amino acid sequence in appropriate positions comprises a number of amino acid residues which can also be found in a foreign T_H epitope then the introduction of a foreign T_H epitope can be accomplished by providing the remaining amino acids of
the foreign epitope by means of amino acid insertion, addition,
10 deletion and substitution. In other words, it is not necessary to introduce a complete T_H epitope by insertion or substitution.

It is preferred that the number of amino acid insertions, deletions, substitutions or additions is at least 2, such as 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20,
15 and 25 insertions, substitutions, additions or deletions. It is furthermore preferred that the number of amino acid insertions, substitutions, additions or deletions is not in excess of 150, such as at most 100, at most 90, at most 80, and at most 70. It is especially preferred that the number of substitutions,
20 insertions, deletions, or additions does not exceed 60, and in particular the number should not exceed 50 or even 40. Most preferred is a number of not more than 30. With respect to amino acid additions, it should be noted that these, when the
resulting construct is in the form of a fusion polypeptide, is
25 often considerably higher than 150.

Preferred embodiments of the invention include modification by introducing at least one foreign immunodominant T_H epitope. It will be understood that the question of immune dominance of a T_H epitope depends on the animal species in question. As used
30 herein, the term "immunodominance" simply refers to epitopes which in the vaccinated individual gives rise to a significant immune response, but it is a well-known fact that a T_H epitope which is immunodominant in one individual is not necessarily immunodominant in another individual of the same species, even

though it may be capable of binding MHC-II molecules in the latter individual.

Another important point is the issue of MHC restriction of T_H epitopes. In general, naturally occurring T_H epitopes are MHC restricted, i.e. a certain peptide constituting a T_H epitope will only bind effectively to a subset of MHC Class II molecules. This in turn has the effect that in most cases the use of one specific T_H epitope will result in a vaccine component which is effective in a fraction of the population only, and depending on the size of that fraction, it can be necessary to include more T_H epitopes in the same molecule, or alternatively prepare a multi-component vaccine wherein the components are GDF-8 variants which are distinguished from each other by the nature of the T_H epitope introduced.

15 If the MHC restriction of the T-cells used is completely unknown (for instance in a situation where the vaccinated animal has a poorly defined MHC composition), the fraction of the animal population covered by a specific vaccine composition can be determined by means of the following formula

20

$$f_{population} = 1 - \prod_{i=1}^n (1 - p_i) \quad (II)$$

-where p_i is the frequency in the population of responders to the i^{th} foreign T-cell epitope present in the vaccine composition, and n is the total number of foreign T-cell epitopes in the vaccine composition. Thus, a vaccine composition containing 3 foreign T-cell epitopes having response frequencies in the population of 0.8, 0.7, and 0.6, respectively, would give

$$1 - 0.2 \times 0.3 \times 0.4 = 0.976$$

-i.e. 97.6 percent of the population will statistically mount an MHC-II mediated response to the vaccine.

The above formula does not apply in situations where a more or less precise MHC restriction pattern of the peptides used is known. If, for instance a certain peptide only binds the human MHC-II molecules encoded by HLA-DR alleles DR1, DR3, DR5, and DR7, then the use of this peptide together with another peptide which binds the remaining MHC-II molecules encoded by HLA-DR alleles will accomplish 100% coverage in the population in question. Likewise, if the second peptide only binds DR3 and DR5, the addition of this peptide will not increase the coverage at all. If one bases the calculation of population response purely on MHC restriction of T-cell epitopes in the vaccine, the fraction of the population covered by a specific vaccine composition can be determined by means of the following formula:

$$f_{\text{population}} = 1 - \prod_{j=1}^3 (1 - \phi_j)^2 \quad (\text{III})$$

-wherein ϕ_j is the sum of frequencies in the population of allelic haplotypes encoding MHC molecules which bind any one of the T-cell epitopes in the vaccine and which belong to the j^{th} of the 3 known HLA loci (DP, DR and DQ); in practice, it is first determined which MHC molecules will recognize each T-cell epitope in the vaccine and thereafter these MHC molecules are listed by type (DP, DR and DQ) - then, the individual frequencies of the different listed allelic haplotypes are summed for each type, thereby yielding ϕ_1 , ϕ_2 , and ϕ_3 .

It may occur that the value p_i in formula II exceeds the corresponding theoretical value π_i :

$$\pi_i = 1 - \prod_{j=1}^3 (1 - v_j)^2 \quad (\text{IV})$$

-wherein v_j is the sum of frequencies in the population of allelic haplotypes encoding MHC molecules which bind the i^{th} T-

cell epitope in the vaccine and which belong to the j^{th} of the 3 known HLA loci (DP, DR and DQ). This means that in $1-\pi_i$ of the population is a frequency of responders of $f_{\text{residual}_i} = (p_i - \pi_i) / (1 - \pi_i)$. Therefore, formula III can be adjusted so as to yield
 5 formula V:

$$f_{\text{population}} = 1 - \prod_{j=1}^3 (1 - \varphi_j)^2 + \left(1 - \prod_{i=1}^n (1 - f_{\text{residual}_i}) \right) \quad (\text{V})$$

-where the term $1 - f_{\text{residual}_i}$ is set to zero if negative. It should be noted that formula V requires that all epitopes have been haplotype mapped against identical sets of haplotypes.

10 Therefore, when selecting T-cell epitopes to be introduced in the GDF-8 analogue, it is for the purposes of optimisation important to include all knowledge of the epitopes which is available: 1) The frequency of responders in the population to each epitope, 2) MHC restriction data, and 3) frequency in the
 15 population of the relevant haplotypes.

There exists a number of naturally occurring "promiscuous" T-cell epitopes which are active in a large proportion of individuals of an animal species or an animal population and these are preferably introduced in the vaccine thereby reducing
 20 the need for a very large number of different GDF-8 analogues in the same vaccine.

The promiscuous epitope can according to the invention be a naturally occurring human T-cell epitope such as epitopes from tetanus toxoid (e.g. the P2 and P30 epitopes), diphtheria
 25 toxoid, Influenza virus hemagglutinin (HA), and *P. falciparum* CS antigen. Of course, when vaccinating other animals than humans care should be taken to utilise naturally occurring T-cell epitopes which are promiscuous in the animal in question.

Over the years a number of other promiscuous T-cell epitopes have been identified. Especially peptides capable of binding a large proportion of HLA-DR molecules encoded by the different HLA-DR alleles have been identified and these are all possible
 5 T-cell epitopes to be introduced in the GDF-8 analogues used according to the present invention. Cf. also the epitopes discussed in the following references which are hereby all incorporated by reference herein: WO 98/23635 (Frazer IH et al., assigned to The University of Queensland); Southwood S et al.,
 10 1998, J. Immunol. **160**: 3363-3373; Sinigaglia F et al., 1988, Nature **336**: 778-780; Chicz RM et al., 1993, J. Exp. Med **178**: 27-47; Hammer J et al., 1993, Cell **74**: 197-203; and Falk K et al., 1994, Immunogenetics **39**: 230-242. The latter reference also deals with HLA-DQ and -DP ligands. All epitopes listed in these
 15 5 references are relevant as candidate natural epitopes to be used in the present invention, as are epitopes which share common motifs with these.

Alternatively, the epitope can be any artificial T-cell epitope which is capable of binding a large proportion of MHC Class II
 20 molecules. In this context the pan DR epitope peptides ("PADRE") described in WO 95/07707 and in the corresponding paper Alexander J et al., 1994, Immunity **1**: 751-761 (both disclosures are incorporated by reference herein) are interesting candidates for epitopes to be used according to the present invention. It
 25 should be noted that the most effective PADRE peptides disclosed in these papers carry D-amino acids in the C- and N-termini in order to improve stability when administered. However, the present invention primarily aims at incorporating the relevant epitopes as part of the modified GDF-8 which should then
 30 subsequently be broken down enzymatically inside the lysosomal compartment of APCs to allow subsequent presentation in the context of an MHC-II molecule and therefore it is not expedient to incorporate D-amino acids in the epitopes used in the present invention.

One especially preferred PADRE peptide is the one having the amino acid sequence AKFVAAWTLKAAA or an immunologically effective subsequence thereof. This, and other epitopes having the same lack of MHC restriction are preferred T-cell epitopes which should be present in the GDF-8 analogues used in the inventive method. Such super-promiscuous epitopes will allow for the most simple embodiments of the invention wherein only one single modified GDF-8 is presented to the vaccinated animal's immune system.

10 As mentioned above, the modification of GDF-8 can also include the introduction of a first moiety which targets the modified GDF-8 to an APC or a B-lymphocyte. For instance, the first moiety can be a specific binding partner for a B-lymphocyte specific surface antigen or for an APC specific surface antigen. 15 Many such specific surface antigens are known in the art. For instance, the moiety can be a carbohydrate for which there is a receptor on the B-lymphocyte or on the APC (e.g. mannan or mannose). Alternatively, the second moiety can be a hapten. Also an antibody fragment which specifically recognizes a surface 20 molecule on APCs or lymphocytes can be used as a first moiety (the surface molecule can e.g. be an FC γ receptor of macrophages and monocytes, such as FC γ RI or, alternatively any other specific surface marker such as CD40 or CTLA-4). It should be noted that all these exemplary targeting molecules can be used 25 as part of an adjuvant also, cf. below.

As an alternative or supplement to targeting the modified GDF-8 polypeptide to a certain cell type in order to achieve an enhanced immune response, it is possible to increase the level of responsiveness of the immune system by including the above- 30 mentioned second moiety which stimulates the immune system. Typical examples of such second moieties are cytokines, and heat-shock proteins, as well as effective parts thereof.

Suitable cytokines to be used according to the invention are those which will normally also function as adjuvants in a vaccine composition, i.e. for instance interferon γ (IFN- γ), Flt3L, interleukin 1 (IL-1), interleukin 2 (IL-2), interleukin 4 (IL-4), interleukin 6 (IL-6), interleukin 12 (IL-12), interleukin 13 (IL-13), interleukin 15 (IL-15), and granulocyte-macrophage colony stimulating factor (GM-CSF); alternatively, the functional part of the cytokine molecule may suffice as the second moiety. With respect to the use of such cytokines as adjuvant substances, cf. the discussion below.

According to the invention, suitable heat shock proteins used as the second moiety can be HSP70, HSP90, HSC70, GRP94 (also known as gp96, cf. Wearsch PA et al. 1998, Biochemistry 37: 5709-19), and CRT (calreticulin).

Alternatively, the second moiety can be a toxin, such as listeriolysin (LLO), lipid A and heat-labile enterotoxin. Also, a number of mycobacterial derivatives such as MDP (muramyl dipeptide) and the trehalose diesters TDM and TDE are interesting possibilities.

Also the possibility of introducing a third moiety which enhances the presentation of the modified GDF-8 to the immune system is an important embodiment of the invention. The art has shown several examples of this principle. For instance, it is known that the palmitoyl lipidation anchor in the *Borrelia burgdorferi* protein OspA can be utilised so as to provide self-adjuvating polypeptides (cf. e.g. WO 96/40718). It seems that the lipidated proteins form up micelle-like structures with a core consisting of the lipidation anchor parts of the polypeptides and the remaining parts of the molecule protruding therefrom, resulting in multiple presentations of the antigenic determinants. Hence, the use of this and related approaches using different lipidation anchors (e.g. a myristyl group, a myristyl group, a farnesyl group, a geranyl-geranyl group, a

GPI-anchor, and an N-acyl diglyceride group) are preferred embodiments of the invention, especially since the provision of such a lipidation anchor in a recombinantly produced protein is fairly straightforward and merely requires use of e.g. a naturally occurring signal sequence as a fusion partner for the modified GDF-8 polypeptide. Another possibility is use of the C3d fragment of complement factor C3 or C3 itself (cf. Dempsey *et al.*, 1996, *Science* **271**, 348-350 and Lou & Kohler, 1998, *Nature Biotechnology* **16**, 458-462).

- 10 An alternative embodiment of the invention which also results in the preferred presentation of multiple (e.g. at least 2) copies of the important epitopic regions of GDF-8 to the immune system is the covalent or non-covalent coupling of GDF-8, subsequence or variants thereof to certain carrier molecules. For instance, polymers can be used, e.g. carbohydrates such as dextran, cf. e.g. Lees A *et al.*, 1994, *Vaccine* **12**: 1160-1166; Lees A *et al.*, 1990, *J Immunol.* **145**: 3594-3600, but also mannose and mannan are useful alternatives. Integral membrane proteins from e.g. *E. coli* and other bacteria are also useful conjugation partners.
- 20 The traditional carrier molecules such as keyhole limpet hemocyanin (KLH), tetanus toxoid, diphtheria toxoid, and bovine serum albumin (BSA) are also preferred and useful conjugation partners.

Certain areas of native GDF-8 are believed to be superiorly suited for performing modifications. It is predicted that modifications in at least one of the following subsequences of C-terminal GDF-8 will result in suitable immunogenic molecules: Residues 18-41, 49-69, or 79-104 in SEQ ID NO: 11 or 12, or corresponding subsequences from GDF-8 polypeptides of different origin than human, bovine, porcine, chicken or turkey. Considerations underlying these chosen areas are a) preservation of B-cell epitopes, b) preservation of secondary, tertiary and quaternary structures etc. At any rate, as discussed herein, it is fairly easy to screen a set of modified GDF-8 molecules

(which have all been subjected to introduction of a T-cell epitope in different locations) for immune reactivity with antibodies raised against a native GDF-8. It is especially preferred that the modification is performed by substituting an amino acid sequence with at least one amino acid sequence of equal or different length which contains a foreign T_H epitope. The rationale behind such constructs is discussed in detail in the examples.

Formulation of GDF-8 and modified GDF-8 polypeptides

- 10 When effecting presentation of the GDF-8 polypeptide or the modified GDF-8 polypeptide to an animal's immune system by means of administration thereof to the animal, the formulation of the polypeptide follows the principles generally acknowledged in the art.
- 15 Preparation of vaccines which contain peptide sequences as active ingredients is generally well understood in the art, as exemplified by U.S. Patents 4,608,251; 4,601,903; 4,599,231; 4,599,230; 4,596,792; and 4,578,770, all incorporated herein by reference. Typically, such vaccines are prepared as injectables
- 20 either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid prior to injection may also be prepared. The preparation may also be emulsified. The active immunogenic ingredient is often mixed with excipients which are pharmaceutically acceptable and compatible with the
- 25 active ingredient. Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol, or the like, and combinations thereof. In addition, if desired, the vaccine may contain minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, or adjuvants which
- 30 enhance the effectiveness of the vaccines; cf. the detailed discussion of adjuvants below.

The vaccines are conventionally administered parenterally, by injection, for example, either subcutaneously, intracutaneously, intradermally, subdermally or intramuscularly. Additional formulations which are suitable for other modes of administration include suppositories and, in some cases, oral, buccal, sublingual, intraperitoneal, intravaginal, anal, epidural, spinal, and intracranial formulations. For ~~suppositories, traditional binders and carriers may include, for~~ example, polyalkylene glycols or triglycerides; such suppositories may be formed from mixtures containing the active ingredient in the range of 0.5% to 10%, preferably 1-2%. Oral formulations include such normally employed excipients as, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, and the like. These compositions take the form of solutions, suspensions, tablets, pills, capsules, sustained release formulations or powders and contain 10-95% of active ingredient, preferably 25-70%. For oral formulations, cholera toxin is an interesting formulation partner (and also a possible conjugation partner).

The polypeptides may be formulated into the vaccine as neutral or salt forms. Pharmaceutically acceptable salts include acid addition salts (formed with the free amino groups of the peptide) and which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like. Salts formed with the free carboxyl groups may also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine, procaine, and the like.

The vaccines are administered in a manner compatible with the dosage formulation, and in such amount as will be therapeutically effective and immunogenic. The quantity to be

administered depends on the subject to be treated, including, e.g., the capacity of the individual's immune system to mount an immune response, and the degree of protection desired. Suitable dosage ranges are of the order of several hundred micrograms active ingredient per vaccination with a preferred range from about 0.1 µg to 2000 µg (even though higher amounts in the 1-10 mg range are contemplated), such as in the range from about 0.5 µg to 1000 µg, preferably in the range from 1 µg to 500 µg and especially in the range from about 10 µg to 100 µg. Suitable regimens for initial administration and booster shots are also variable but are typified by an initial administration followed by subsequent inoculations or other administrations.

The manner of application may be varied widely. Any of the conventional methods for administration of a vaccine are applicable. These include oral application on a solid physiologically acceptable base or in a physiologically acceptable dispersion, parenterally, by injection or the like. The dosage of the vaccine will depend on the route of administration and will vary according to the age of the person to be vaccinated and the formulation of the antigen.

Some of the polypeptides of the vaccine are sufficiently immunogenic in a vaccine, but for some of the others the immune response will be enhanced if the vaccine further comprises an adjuvant substance.

Various methods of achieving adjuvant effect for the vaccine are known. General principles and methods are detailed in "The Theory and Practical Application of Adjuvants", 1995, Duncan E.S. Stewart-Tull (ed.), John Wiley & Sons Ltd, ISBN 0-471-95170-6, and also in "Vaccines: New Generation Immunological Adjuvants", 1995, Gregoriadis G et al. (eds.), Plenum Press, New York, ISBN 0-306-45283-9, both of which are hereby incorporated by reference herein.

It is especially preferred to use an adjuvant which can be demonstrated to facilitate breaking of the autotolerance to autoantigens; in fact, this is essential in cases where unmodified GDF-8 is used as the active ingredient in the
 5 autovaccine. Non-limiting examples of suitable adjuvants are selected from the group consisting of an immune targeting adjuvant; an immune modulating adjuvant such as a toxin, a cytokine, and a mycobacterial derivative; an oil formulation; a
 10 polymer; a micelle forming adjuvant; a saponin; an immunostimulating complex matrix (ISCOM matrix); a particle; DDA; aluminium adjuvants; DNA adjuvants; γ -inulin; and an encapsulating adjuvant. In general it should be noted that the disclosures above which relate to compounds and agents useful as first, second and third moieties in the analogues also refer
 15 *mutatis mutandis* to their use in the adjuvant of a vaccine of the invention.

The application of adjuvants include use of agents such as aluminum hydroxide or phosphate (alum), commonly used as 0.05 to 0.1 percent solution in buffered saline, admixture with
 20 synthetic polymers of sugars (e.g. Carbopol®) used as 0.25 percent solution, aggregation of the protein in the vaccine by heat treatment with temperatures ranging between 70° to 101°C for 30 second to 2 minute periods respectively and also aggregation by means of cross-linking agents are possible.
 25 Aggregation by reactivation with pepsin treated antibodies (Fab fragments) to albumin, mixture with bacterial cells such as *C. parvum* or endotoxins or lipopolysaccharide components of gram-negative bacteria, emulsion in physiologically acceptable oil vehicles such as mannide mono-oleate (Aracel A) or emulsion with
 30 20 percent solution of a perfluorocarbon (Fluosol-DA) used as a block substitute may also be employed. Admixture with oils such as squalene and IFA is also preferred.

According to the invention DDA (dimethyldioctadecylammonium bromide) is an interesting candidate for an adjuvant as is DNA

and γ -inulin, but also Freund's complete and incomplete adjuvants as well as *quillaja* saponins such as QuilA and QS21 are interesting as is RIBI. Further possibilities are monophosphoryl lipid A (MPL), the above mentioned C3 and C3d, 5 and muramyl dipeptide (MDP).

Liposome formulations are also known to confer adjuvant effects, and therefore liposome adjuvants are preferred according to the invention.

Also immunostimulating complex matrix type (ISCOM® matrix) 10 adjuvants are preferred choices according to the invention, especially since it has been shown that this type of adjuvants are capable of up-regulating MHC Class II expression by APCs. An ISCOM® matrix consists of (optionally fractionated) saponins (triterpenoids) from *Quillaja saponaria*, cholesterol, and 15 phospholipid. When admixed with the immunogenic protein, the resulting particulate formulation is what is known as an ISCOM particle where the saponin constitutes 60-70% w/w, the cholesterol and phospholipid 10-15% w/w, and the protein 10-15% w/w. Details relating to composition and use of 20 immunostimulating complexes can e.g. be found in the above-mentioned text-books dealing with adjuvants, but also Morein B et al., 1995, Clin. Immunother. 3: 461-475 as well as Barr IG and Mitchell GF, 1996, Immunol. and Cell Biol. 74: 8-25 (both incorporated by reference herein) provide useful instructions 25 for the preparation of complete immunostimulating complexes.

Another highly interesting (and thus, preferred) possibility of achieving adjuvant effect is to employ the technique described in Gosselin et al., 1992 (which is hereby incorporated by reference herein). In brief, the presentation of a relevant 30 antigen such as an antigen of the present invention can be enhanced by conjugating the antigen to antibodies (or antigen binding antibody fragments) against the Fc γ receptors on monocytes/macrophages. Especially conjugates between antigen and

anti-FcγRI have been demonstrated to enhance immunogenicity for the purposes of vaccination.

Other possibilities involve the use of the targeting and immune modulating substances (i.a. cytokines) mentioned above as
5 candidates for the first and second moieties in the modified versions of GDF-8. In this connection, also synthetic inducers of cytokines like poly I:C are possibilities.

Suitable mycobacterial derivatives are selected from the group consisting of muramyl dipeptide, complete Freund's adjuvant,
10 RIBI, and a diester of trehalose such as TDM and TDE.

Suitable immune targeting adjuvants are selected from the group consisting of CD40 ligand and CD40 antibodies or specifically binding fragments thereof (cf. the discussion above), mannose, a Fab fragment, and CTLA-4.

15 Suitable polymer adjuvants are selected from the group consisting of a carbohydrate such as dextran, PEG, starch, mannan, and mannose; a plastic polymer such as; and latex such as latex beads.

Yet another interesting way of modulating an immune response is
20 to include the GDF-8 immunogen (optionally together with adjuvants and pharmaceutically acceptable carriers and vehicles) in a "virtual lymph node" (VLN) (a proprietary medical device developed by ImmunoTherapy, Inc., 360 Lexington Avenue, New York, NY 10017-6501). The VLN (a thin tubular device) mimics the
25 structure and function of a lymph node. Insertion of a VLN under the skin creates a site of sterile inflammation with an upsurge of cytokines and chemokines. T- and B-cells as well as APCs rapidly respond to the danger signals, home to the inflamed site and accumulate inside the porous matrix of the VLN. It has been
30 shown that the necessary antigen dose required to mount an immune response to an antigen is reduced when using the VLN and

that immune protection conferred by vaccination using a VLN surpassed conventional immunization using Ribi as an adjuvant. The technology is *i.a.* described briefly in Gelber C et al., 1998, "Elicitation of Robust Cellular and Humoral Immune Responses to Small Amounts of Immunogens Using a Novel Medical Device Designated the Virtual Lymph Node", in: "From the Laboratory to the Clinic, Book of Abstracts, October 12th - 15th 1998, Seascape Resort, Aptos, California".

It is expected that the vaccine should be administered at least once a year, such as at least 1, 2, 3, 4, 5, 6, and 12 times a year. More specifically, 1-12 times per year is expected, such as 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12 times a year to an individual in need thereof. It has previously been shown that the memory immunity induced by the use of the preferred autovaccines according to the invention is not permanent, and therefor the immune system needs to be periodically challenged with the analogues.

Due to genetic variation, different individuals may react with immune responses of varying strength to the same polypeptide. Therefore, the vaccine according to the invention may comprise several different polypeptides in order to increase the immune response, cf. also the discussion above concerning the choice of foreign T-cell epitope introductions. The vaccine may comprise two or more polypeptides, where all of the polypeptides are as defined above.

The vaccine may consequently comprise 3-20 different modified or unmodified polypeptides, such as 3-10 different polypeptides. However, normally the number of polypeptides will be sought kept to a minimum such as 1 or 2 polypeptides.

Nucleic acid vaccination

As an alternative to classic administration of a peptide-based vaccine, the technology of nucleic acid vaccination (also known as "nucleic acid immunisation", "genetic immunisation", and
5 "gene immunisation") offers a number of attractive features.

First, in contrast to the traditional vaccine approach, nucleic acid vaccination does not require resource consuming large-scale production of the immunogenic agent (e.g. in the form of industrial scale fermentation of microorganisms producing
10 modified GDF-8). Furthermore, there is no need to devise purification and refolding schemes for the immunogen. And finally, since nucleic acid vaccination relies on the biochemical apparatus of the vaccinated individual in order to produce the expression product of the nucleic acid introduced,
15 the optimum posttranslational processing of the expression product is expected to occur; this is especially important in the case of autovaccination, since, as mentioned above, a significant fraction of the original GDF-8 B-cell epitopes should be preserved in the modified molecule, and since B-cell
20 epitopes in principle can be constituted by parts of any (bio)molecule (e.g. carbohydrate, lipid, protein etc.). Therefore, native glycosylation and lipidation patterns of the immunogen may very well be of importance for the overall immunogenicity and this is expected to be ensured by having the
25 host producing the immunogen.

Hence, a preferred embodiment of the invention comprises effecting presentation of modified GDF-8 to the immune system by introducing nucleic acid(s) encoding the modified GDF-8 into the animal's cells and thereby obtaining *in vivo* expression by the
30 cells of the nucleic acid(s) introduced.

In this embodiment, the introduced nucleic acid is preferably DNA which can be in the form of naked DNA, DNA formulated with

charged or uncharged lipids, DNA formulated in liposomes, DNA included in a viral vector, DNA formulated with a transfection-facilitating protein or polypeptide, DNA formulated with a targeting protein or polypeptide, DNA formulated with Calcium precipitating agents, DNA coupled to an inert carrier molecule, and DNA formulated with an adjuvant. In this context it is noted that practically all considerations pertaining to the use of adjuvants in traditional vaccine formulation apply for the formulation of DNA vaccines. Hence, all disclosures herein which relate to use of adjuvants in the context of polypeptide based vaccines apply *mutatis mutandis* to their use in nucleic acid vaccination technology.

As for routes of administration and administration schemes of polypeptide based vaccines which have been detailed above, these are also applicable for the nucleic acid vaccines of the invention and all discussions above pertaining to routes of administration and administration schemes for polypeptides apply *mutatis mutandis* to nucleic acids. To this should be added that nucleic acid vaccines can suitably be administered intravenously and intraarterially. Furthermore, it is well-known in the art that nucleic acid vaccines can be administered by use of a so-called gene gun, and hence also this and equivalent modes of administration are regarded as part of the present invention. Finally, also the use of a VLN in the administration of nucleic acids has been reported to yield good results, and therefore this particular mode of administration is particularly preferred.

Furthermore, the nucleic acid(s) used as an immunization agent can contain regions encoding the 1st, 2nd and/or 3rd moieties, e.g. in the form of the immunomodulating substances described above such as the cytokines discussed as useful adjuvants. A preferred version of this embodiment encompasses having the coding region for the analogue and the coding region for the immunomodulator in different reading frames or at least under

the control of different promoters. Thereby it is avoided that the analogue or epitope is produced as a fusion partner to the immunomodulator. Alternatively, two distinct nucleotide fragments can be used, but this is less preferred because of the
 5 advantage of ensured co-expression when having both coding regions included in the same molecule.

Accordingly, the invention also relates to a composition for inducing production of antibodies against GDF-8, the composition comprising

- 10 - a nucleic acid fragment or a vector of the invention (cf. the discussion of vectors below), and
- a pharmaceutically and immunologically acceptable vehicle and/or carrier and/or adjuvant as discussed above.

Under normal circumstances, the GDF-8 variant-encoding nucleic
 15 acid is introduced in the form of a vector wherein expression is under control of a viral promoter. For more detailed discussions of vectors and DNA fragments according to the invention, cf. the discussion below. Also, detailed disclosures relating to the formulation and use of nucleic acid vaccines are available, cf.
 20 Donnelly JJ et al, 1997, Annu. Rev. Immunol. **15**: 617-648 and Donnelly JJ et al., 1997, Life Sciences **60**: 163-172. Both of these references are incorporated by reference herein.

Live vaccines

A third alternative for effecting presentation of modified GDF-8
 • 25 to the immune system is the use of live vaccine technology. In live vaccination, presentation to the immune system is effected by administering, to the animal, a non-pathogenic microorganism which has been transformed with a nucleic acid fragment encoding a modified GDF-8 or with a vector incorporating such a nucleic
 30 acid fragment. The non-pathogenic microorganism can be any suitable attenuated bacterial strain (attenuated by means of passaging or by means of removal of pathogenic expression

products by recombinant DNA technology), e.g. *Mycobacterium bovis* BCG., non-pathogenic *Streptococcus* spp., *E. coli*, *Salmonella* spp., *Vibrio cholerae*, *Shigella*, etc. Reviews dealing with preparation of state-of-the-art live vaccines can e.g. be found in Saliou P, 1995, Rev. Prat. 45: 1492-1496 and Walker PD, 1992, Vaccine 10: 977-990, both incorporated by reference herein. For details about the nucleic acid fragments and vectors used in such live vaccines, cf. the discussion below.

As an alternative to bacterial live vaccines, the nucleic acid fragment of the invention discussed below can be incorporated in a non-virulent viral vaccine vector such as a vaccinia strain or any other suitable pox virus which will be infectious in the animal to be vaccinated.

Normally, the non-pathogenic microorganism or virus is administered only once to the animal, but in certain cases it may be necessary to administer the microorganism more than once in a lifetime in order to maintain protective immunity. It is even contemplated that immunization schemes as those detailed above for polypeptide vaccination will be useful when using live or virus vaccines.

Alternatively, live or virus vaccination is combined with previous or subsequent polypeptide and/or nucleic acid vaccination. For instance, it is possible to effect primary immunization with a live or virus vaccine followed by subsequent booster immunizations using the polypeptide or nucleic acid approach.

The microorganism or virus can be transformed with nucleic acid(s) containing regions encoding the 1st, 2nd and/or 3rd moieties, e.g. in the form of the immunomodulating substances described above such as the cytokines discussed as useful adjuvants. A preferred version of this embodiment encompasses having the coding region for the analogue and the coding region

for the immunomodulator in different reading frames or at least under the control of different promoters. Thereby it is avoided that the analogue or epitopes are produced as fusion partners to the immunomodulator. Alternatively, two distinct nucleotide
5 fragments can be used as transforming agents. Of course, having the 1st and/or 2nd and/or 3rd moieties in the same reading frame can provide as an expression product, an analogue of the invention, and such an embodiment is especially preferred
according to the present invention.

10 Use of the method of the invention in meat production and in disease treatment

As will be appreciated from the discussions above, the provision of the method of the invention for down-regulation of GDF-8 activity allows for stimulation of the growth of skeletal muscle
15 mass in animals. Hence, an important embodiment of the method of the invention for down-regulating GDF-8 activity comprises increasing the skeletal muscle mass of an animal, the method comprising down-regulating GDF-8 activity according to the method of the invention to such an extent that the muscle mass
20 is statistically significantly increased (at a confidence level of 95%) and with at least 5%, when compared to control animals exhibiting a normal GDF-8 activity.

The increase in muscle mass may preferably be higher, such as at least 10, 15, 20, 25, 30, 35, 40, or even 45%, cf. the increases
25 in muscle mass which have been observed in transgenic mice and naturally occurring GDF-8 deficient animals.

The muscle mass can be determined by any convenient method known in the art for assessing total and/or relative muscle mass.

30 An anti-GDF-8 vaccine could potentially also be useful for treatment of certain human diseases such as cancer cachexia, where muscular atrophy is a pronounced phenomenon, and it is

also a feasible means of treatment/amelioration in other atrophic muscular diseases.

Peptides, polypeptides, and compositions of the invention

As will be apparent from the above, the present invention is
5 based on the concept of immunising individuals against the GDF-8 antigen in order to obtain an increase in muscle growth rate.

The preferred way of obtaining such an immunization is to use modified versions of GDF-8, thereby providing molecules which have not previously been disclosed in the art.

10 It is believed that the modified GDF-8 molecules discussed herein are inventive in their own right, and therefore an important part of the invention pertains to a GDF-8 analogue which is derived from an animal GDF-8 wherein is introduced a modification which has as a result that immunization of the
15 animal with the analogue induces production of antibodies reacting specifically with the unmodified GDF-8 polypeptide. Preferably, the nature of the modification conforms with the types of modifications described above when discussing various embodiments of the method of the invention when using modified
20 GDF-8. Hence, any disclosure presented herein pertaining to modified GDF-8 molecules are relevant for the purpose of describing the GDF-8 analogues of the invention, and any such disclosures apply *mutatis mutandis* to the description of these analogues.

25 It should be noted that preferred modified GDF-8 molecules comprise modifications which results in a polypeptide having a sequence identity of at least 70% with GDF-8 or with a subsequence thereof of at least 10 amino acids in length. Higher sequence identities are preferred, e.g. at least 75% or even at
30 least 80% or 85%. The sequence identity for proteins and nucleic acids can be calculated as $(N_{ref} - N_{dif}) \cdot 100 / N_{ref}$, wherein N_{dif} is the total number of non-identical residues in the two sequences

when aligned and wherein N_{ref} is the number of residues in one of the sequences. Hence, the DNA sequence AGTCAGTC will have a sequence identity of 75% with the sequence AATCAATC ($N_{dif}=2$ and $N_{ref}=8$).

5 The invention also pertains to compositions useful in exercising the method of the invention. Hence, the invention also relates to an immunogenic composition comprising an immunogenically effective amount of a GDF-8 polypeptide which is a self-protein in an animal, said GDF-8 polypeptide being formulated together
10 with an immunologically acceptable adjuvant so as to break the animal's autotolerance towards the GDF-8 polypeptide, the composition further comprising a pharmaceutically and immunologically acceptable vehicle and/or carrier. In other words, this part of the invention pertains to the formulations
15 of naturally occurring GDF-8 polypeptides which have been described in connection with embodiments of the method of the invention.

The invention also relates to an immunogenic composition comprising an immunologically effective amount of a GDF-8
20 analogue defined above, said composition further comprising a pharmaceutically and immunologically acceptable diluent and/or vehicle and/or carrier and/or excipient and optionally an adjuvant. In other words, this part of the invention concerns formulations of modified GDF-8, essentially as described
25 hereinabove. The choice of adjuvants, carriers, and vehicles is accordingly in line with what has been discussed above when referring to formulation of modified and unmodified GDF-8 for use in the inventive method for the down-regulation of GDF-8.

The polypeptides are prepared according to methods well-known in
30 the art. Longer polypeptides are normally prepared by means of recombinant gene technology including introduction of a nucleic acid sequence encoding the GDF-8 analogue into a suitable vector, transformation of a suitable host cell with the vector,

expression of the nucleic acid sequence, recovery of the expression product from the host cells or their culture supernatant, and subsequent purification and optional further modification, e.g. refolding or derivatisation.

- 5 Shorter peptides are preferably prepared by means of the well-known techniques of solid- or liquid-phase peptide synthesis. However, recent advances in this technology has rendered possible the production of full-length polypeptides and proteins by these means, and therefore it is also within the scope of the
10 present invention to prepare the long constructs by synthetic means.

Nucleic acid fragments and vectors of the invention

It will be appreciated from the above disclosure that modified GDF-8 polypeptides can be prepared by means of recombinant gene
15 technology but also by means of chemical synthesis or semisynthesis; the latter two options are especially relevant when the modification consists in coupling to protein carriers (such as KLH, diphtheria toxoid, tetanus toxoid, and BSA) and non-proteinaceous molecules such as carbohydrate polymers and of
20 course also when the modification comprises addition of side chains or side groups to a GDF-8 polypeptide-derived peptide chain.

For the purpose of recombinant gene technology, and of course also for the purpose of nucleic acid immunization, nucleic acid
25 fragments encoding modified GDF-8 are important chemical products. Hence, an important part of the invention pertains to a nucleic acid fragment which encodes a GDF-8 analogue, i.e. a GDF-8 derived polypeptide which either comprises the natural GDF-8 sequence to which has been added or inserted a fusion
30 partner or, preferably a GDF-8 derived polypeptide wherein has been introduced a foreign T-cell epitope by means of insertion and/or addition, preferably by means of substitution and/or

deletion. The nucleic acid fragments of the invention are either DNA or RNA fragments.

The nucleic acid fragments of the invention will normally be inserted in suitable vectors to form cloning or expression
5 vectors carrying the nucleic acid fragments of the invention; such novel vectors are also part of the invention. Details concerning the construction of these vectors of the invention

will be discussed in context of transformed cells and microorganisms below. The vectors can, depending on purpose and
10 type of application, be in the form of plasmids, phages, cosmids, mini-chromosomes, or virus, but also naked DNA which is only expressed transiently in certain cells is an important vector. Preferred cloning and expression vectors of the invention are capable of autonomous replication, thereby
15 enabling high copy-numbers for the purposes of high-level expression or high-level replication for subsequent cloning.

The general outline of a vector of the invention comprises the following features in the 5'-3' direction and in operable linkage: a promoter for driving expression of the nucleic acid
20 fragment of the invention, optionally a nucleic acid sequence encoding a leader peptide enabling secretion (to the extracellular phase or, where applicable, into the periplasma of a bacterium) of or integration into the membrane of the polypeptide fragment, the nucleic acid fragment of the
25 invention, and optionally a nucleic acid sequence encoding a terminator. When operating with expression vectors in producer strains or cell-lines it is for the purposes of genetic stability of the transformed cell preferred that the vector when introduced into a host cell is integrated in the host cell
30 genome. In contrast, when working with vectors to be used for effecting *in vivo* expression in an animal (*i.e.* when using the vector in DNA vaccination) it is for security reasons preferred that the vector is not incapable of being integrated in the host cell genome; typically, naked DNA or non-integrating viral

vectors are used, the choices of which are well-known to the person skilled in the art.

The vectors of the invention are used to transform host cells to produce the modified GDF-8 polypeptide of the invention. Such
5 transformed cells, which are also part of the invention, can be cultured cells or cell lines used for propagation of the nucleic acid fragments and vectors of the invention, or used for recombinant production of the modified GDF-8 polypeptides of the invention. Alternatively, the transformed cells can be suitable
10 live vaccine strains wherein the nucleic acid fragment (one single or multiple copies) have been inserted so as to effect secretion or integration into the bacterial membrane or cell-wall of the modified GDF-8.

Preferred transformed cells of the invention are microorganisms
15 such as bacteria (such as the species *Escherichia* [e.g. *E.coli*], *Bacillus* [e.g. *Bacillus subtilis*], *Salmonella*, or *Mycobacterium* [preferably non-pathogenic, e.g. *M. bovis* BCG]), yeasts (such as *Saccharomyces cerevisiae*), and protozoans. Alternatively, the transformed cells are derived from a multicellular organism such
20 as a fungus, an insect cell, a plant cell, or a mammalian cell. Recent results have shown great promise in the use of a commercially available *Drosophila melanogaster* cell line (the Schneider 2 (S₂) cell line and vector system available from Invitrogen) for the recombinant production of IL-5 analogues of
25 the invention, and therefore this expression system is particularly preferred, also for the purposes of the present invention.

For the purposes of cloning and/or optimized expression it is preferred that the transformed cell is capable of replicating
30 the nucleic acid fragment of the invention. Cells expressing the nucleic acid fragment are preferred useful embodiments of the invention; they can be used for small-scale or large-scale

preparation of the modified GDF-8 or, in the case of non-pathogenic bacteria, as vaccine constituents in a live vaccine.

When producing the modified GDF-8 of the invention by means of transformed cells, it is convenient, although far from essential, that the expression product is either exported out into the culture medium or carried on the surface of the transformed cell.

When an effective producer cell has been identified it is preferred, on the basis thereof, to establish a stable cell line which carries the vector of the invention and which expresses the nucleic acid fragment encoding the modified GDF-8. Preferably, this stable cell line secretes or carries the GDF-8 analogue of the invention, thereby facilitating purification thereof.

In general, plasmid vectors containing replicon and control sequences which are derived from species compatible with the host cell are used in connection with the hosts. The vector ordinarily carries a replication site, as well as marking sequences which are capable of providing phenotypic selection in transformed cells. For example, *E. coli* is typically transformed using pBR322, a plasmid derived from an *E. coli* species (see, e.g., Bolivar et al., 1977). The pBR322 plasmid contains genes for ampicillin and tetracycline resistance and thus provides easy means for identifying transformed cells. The pBR plasmid, or other microbial plasmid or phage must also contain, or be modified to contain, promoters which can be used by the prokaryotic microorganism for expression.

Those promoters most commonly used in recombinant DNA construction include the B-lactamase (penicillinase) and lactose promoter systems (Chang et al., 1978; Itakura et al., 1977; Goeddel et al., 1979) and a tryptophan (*trp*) promoter system (Goeddel et al., 1979; EP-A-0 036 776). While these are the most

commonly used, other microbial promoters have been discovered and utilized, and details concerning their nucleotide sequences have been published, enabling a skilled worker to ligate them functionally with plasmid vectors (Siebwenlist et al., 1980).

- 5 Certain genes from prokaryotes may be expressed efficiently in *E. coli* from their own promoter sequences, precluding the need for addition of another promoter by artificial means.

In addition to prokaryotes, eukaryotic microbes, such as yeast cultures may also be used, and here the promoter should be
10 capable of driving expression. *Saccharomyces cerevisiae*, or common baker's yeast is the most commonly used among eukaryotic microorganisms, although a number of other strains are commonly available. For expression in *Saccharomyces*, the plasmid YRp7, for example, is commonly used (Stinchcomb et al., 1979; Kingsman
15 et al., 1979; Tschemper et al., 1980). This plasmid already contains the *trp1* gene which provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan for example ATCC No. 44076 or PEP4-1 (Jones, 1977). The presence of the *trp1* lesion as a characteristic of the yeast host cell
20 genome then provides an effective environment for detecting transformation by growth in the absence of tryptophan.

Suitable promoting sequences in yeast vectors include the promoters for 3-phosphoglycerate kinase (Hitzman et al., 1980) or other glycolytic enzymes (Hess et al., 1968; Holland et al.,
25 1978), such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase. In
30 constructing suitable expression plasmids, the termination sequences associated with these genes are also ligated into the expression vector 3' of the sequence desired to be expressed to provide polyadenylation of the mRNA and termination.

Other promoters, which have the additional advantage of transcription controlled by growth conditions are the promoter region for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, and the aforementioned glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose utilization. Any plasmid vector containing a yeast-compatible promoter, origin of replication and termination sequences is suitable.

- 10 In addition to microorganisms, cultures of cells derived from multicellular organisms may also be used as hosts. In principle, any such cell culture is workable, whether from vertebrate or invertebrate culture. However, interest has been greatest in vertebrate cells, and propagation of vertebrate in culture
- 15 (tissue culture) has become a routine procedure in recent years (Tissue Culture, 1973). Examples of such useful host cell lines are VERO and HeLa cells, Chinese hamster ovary (CHO) cell lines, and W138, BHK, COS-7 293, *Drosophila melanogaster* and *Spodoptera frugiperda* (SF) cells (commercially available as complete
- 20 expression systems from i.a. Protein Sciences, 1000 Research Parkway, Meriden, CT 06450, U.S.A. and from Invitrogen), and MDCK cell lines. In the present invention, especially preferred cell lines are S₂ and SF₂₁ available from Invitrogen, PO Box 2312, 9704 CH Groningen, The Netherlands.
- 25 Expression vectors for such cells ordinarily include (if necessary) an origin of replication, a promoter located in front of the gene to be expressed, along with any necessary ribosome binding sites, RNA splice sites, polyadenylation site, and transcriptional terminator sequences.
- 30 For use in mammalian cells, the control functions on the expression vectors are often provided by viral material. For example, commonly used promoters are derived from polyoma, Adenovirus 2, and most frequently Simian Virus 40 (SV40). The

early and late promoters of SV40 virus are particularly useful because both are obtained easily from the virus as a fragment which also contains the SV40 viral origin of replication (Fiers et al., 1978). Smaller or larger SV40 fragments may also be
5 used, provided there is included the approximately 250 bp sequence extending from the *HindIII* site toward the *BglI* site located in the viral origin of replication. Further, it is also possible, and often desirable, to utilize promoter or control
sequences normally associated with the desired gene sequence,
10 provided such control sequences are compatible with the host cell systems.

An origin of replication may be provided either by construction of the vector to include an exogenous origin, such as may be derived from SV40 or other viral (e.g., Polyoma, Adeno, VSV,
15 BPV) or may be provided by the host cell chromosomal replication mechanism. If the vector is integrated into the host cell chromosome, the latter is often sufficient.

Identification of useful GDF-8 analogues

It will be clear to the skilled person that not all variants or
20 modifications of native GDF-8 will have the ability to elicit antibodies in an animal which are cross-reactive with the native form. It is, however, not difficult to set up an effective standard screen for modified GDF-8 molecules which fulfill the minimum requirements for immunological reactivity discussed
25 herein. Hence, another part of the invention concerns a method for the identification of a modified GDF-8 polypeptide which is capable of inducing antibodies against unmodified GDF-8 in an animal species where the unmodified GDF-8 polypeptide is a self-protein, the method comprising

30 - preparing, by means of peptide synthesis or by genetic engineering methods, a set of mutually distinct modified GDF-8 polypeptides wherein amino acids have been added to,

- inserted in, deleted from, or substituted into the amino acid sequence of a GDF-8 polypeptide of the animal species thereby giving rise to amino acid sequences in the set which comprise T-cell epitopes which are foreign to the animal species,
- 5 - testing members of the set for their ability to induce production of antibodies by the animal species against the unmodified GDF-8, and
-
- 10 - isolating the member(s) of the set which significantly induces antibody production against unmodified GDF-8 in the animal species.

In this context, the "set of mutually distinct modified GDF-8 polypeptides" is a collection of non-identical modified GDF-8 polypeptides which have e.g. been selected on the basis of the

15 criteria discussed above (e.g. in combination with studies of circular dichroism, NMR spectra, and/or X-ray diffraction patterns). The set may consist of only a few members but it is contemplated that the set may contain several hundred members.

The test of members of the set can ultimately be performed *in vivo*, but a number of *in vitro* tests can be applied which narrow

20 down the number of modified molecules which will serve the purpose of the invention.

Since the goal of introducing the foreign T-cell epitopes is to support the B-cell response by T-cell help, a prerequisite is

25 that T-cell proliferation is induced by the modified GDF-8. T-cell proliferation can be tested by standardized proliferation assays *in vitro*. In short, a sample enriched for T-cells is obtained from a subject and subsequently kept in culture. The cultured T-cells are contacted with APCs of the subject which

30 have previously taken up the modified molecule and processed it to present its T-cell epitopes. The proliferation of T-cells is monitored and compared to a suitable control (e.g. T-cells in culture contacted with APCs which have processed intact, native

GDF-8). Alternatively, proliferation can be measured by determining the concentration of relevant cytokines released by the T-cells in response to their recognition of foreign T-cells.

Having rendered highly probable that at least one modified GDF-8
5 of the set is capable of inducing antibody production against GDF-8, it is possible to prepare an immunogenic composition comprising at least one modified GDF-8 polypeptide which is
capable of inducing antibodies against unmodified GDF-8 in an
animal species where the unmodified GDF-8 polypeptide is a self-
10 protein, the method comprising admixing the member(s) of the set which significantly induces production of antibodies in the
animal species which are reactive with GDF-8 with a
pharmaceutically and immunologically acceptable carrier and/or
vehicle and/or diluent and/or excipient, optionally in
15 combination with at least one pharmaceutically and
immunologically acceptable adjuvant.

The above aspects of the invention are conveniently carried out by initially preparing a number of mutually distinct nucleic acid sequences or vectors of the invention, inserting these into
20 appropriate expression vectors, transforming suitable host cells with the vectors, and expressing the nucleic acid sequences of the invention. These steps can be followed by isolation of the expression products. It is preferred that the nucleic acid sequences and/or vectors are prepared by methods comprising
25 exercise of a molecular amplification technique such as PCR or by means of nucleic acid synthesis.

PREAMBLE TO EXAMPLES

Expression of the 109 amino acid residue C-terminal region of GDF-8 fused to an N-terminal His-tag has been obtained in *E.*
30 *coli* and the purified fusion protein has been used for immunisation of rabbits. Full-length GDF-8 has been expressed in CHO cells and shown to be secreted as dimers of unprocessed and

processed GDF-8, respectively (McPherron et al., Nature 387, 83-90, 1997). Thus, no problems are expected with expression of the below-discussed GDF-8 autovaccine constructs. The most likely expression systems to be used for production of the GDF-8 AutoVac constructs are *E. coli*, the yeast *P. pastoris* and CHO cells and insect cells such as the above mentioned *Drosophila* cells.

EXAMPLE 1

Vaccine design

10 The rationale behind the present invention is to substitute stretches of amino acid residues in the target protein with foreign or artificial T-cell epitopes e.g. the promiscuous tetanus toxin T-cell epitopes P2 and P30. Preferably these substitutions should only minimally disturb the authentic three
15 dimensional structure of the target protein.

The target protein here is the 109 amino acid residue C-terminal region of GDF-8, the homodimer of which is expected to be the biologically active form of GDF-8. The three-dimensional structure of this region of GDF-8 is not known but based on the
20 structure of the homologous TGF- β protein the model of monomeric GDF-8 shown in Fig. 2 can be anticipated to be reasonably close to reality. In this model of the wild type GDF-8 (wt) α -helices are shown as cylinders and β -sheets are shown as arrows. The cysteine-residues and thus the disulfide-bonds are very closely
25 positioned in the structure.

It should be borne in mind that the presence of the relatively high number cysteins (9) and thus disulfide-bonds in the 109 amino acid residue C-terminal region of GDF-8 limits the possible sites where the foreign T-cell epitopes can be
30 positioned.

In addition to producing the 109 amino acid residue C-terminal region of GDF-8 without substitutions (i.e. residues 267-375 of SEQ ID NOS: 1-10), the following GDF-8 AutoVac constructs are proposed.

5 **GDF-8 P2-1** (SEQ ID NO: 15) is the 109 amino acid residue C-terminal region of GDF-8 with amino acid residues 18-32 substituted by P2.

GDF-8 P2-2 (SEQ ID NO: 16) is the 109 amino acid residue C-terminal region of GDF-8 with amino acid residues 52-66 substituted by P2.

GDF-8 P2-3 (SEQ ID NO: 17) is the 109 amino acid residue C-terminal region of GDF-8 with amino acid residues 83-97 substituted by P2.

15 **GDF-8 P30-1** (SEQ ID NO: 18) is the 109 amino acid residue C-terminal region of GDF-8 with amino acid residues 21-41 substituted by P30.

GDF-8 P30-2 (SEQ ID NO: 19) is the 109 amino acid residue C-terminal region of GDF-8 with amino acid residues 49-69 substituted by P30.

20 **GDF-8 P30-3A** (SEQ ID NO: 20) is the 109 amino acid residue C-terminal region of GDF-8 with amino acid residues 79-99 substituted by P30.

GDF-8 P30-3B (SEQ ID NO: 21) is the 109 amino acid residue C-terminal region of GDF-8 with amino acid residues 84-104 substituted by P30.

GDF-8 dimer (SEQ ID NO: 22) is two copies of the 109 amino acid residue C-terminal region of GDF-8 covalently connected through the P2 and P30 epitopes. In other words the molecule is

comprised of two halves. The first half is the 109 amino acid residue C-terminal region of GDF-8 having P2 fused to its C-terminus while the second half is the 109 amino acid residue C-terminal region of GDF-8 having P30 fused to its N-terminus.

- 5 **GDF-8 ext** (SEQ ID NO: 23) consists of the C-terminal 160 amino acid residue of GDF-8 with residues 16-36 substituted by P30 and residues 37-51 substituted by P2. This construct is the 109 amino acid residue C-terminal region of GDF-8 with an N-terminal extension containing both the P2 and the P30 epitopes.
- 10 In all exemplary constructs except from the latter 2, it is contemplated to produce a variant where Cys73 is substituted with Ser to avoid dimerisation through disulfide-bond formation. In **GDF-8 ext** it is contemplated to perform a similar substitution in the corresponding position (i.e. Cys124 -
15 Ser124).

EXAMPLE 2

In vitro models

- It is contemplated to initially immunise mice with purified GDF-8 variants as described above. After e.g. three immunisations
- 20 antibodies will be measured in ELISA using the non-modified GDF-8 molecule as the antigen. The primary reason for these initial experiments is to confirm that the AutoVac™ technology is applicable for creating anti-GDF-8 cross reactive autoantibodies and, importantly, to identify the optimal dosing and
 - 25 immunisation regimen. It would, however, be very surprising if antibodies against GDF-8 were not raised using the present technology, since the fundamental immunological mechanisms of such a response are most likely to be identical to those which have already been observed for TNF α , cf. WO 98/46642 and WO
30 95/05849.

EXAMPLE 3

In vivo models

Groups of mice will be immunised with the various constructs described in Example 1. The mice will be immunised at the age of 5 about 4 weeks since they have to be immune competent in order to respond to the vaccine. Freund's Complete adjuvant will be used, but experiments will also be performed using an alum adjuvant such as Adjuphos™, which has previously been used successfully in admixture with TNF α autovaccine constructs. Adjuphos™ is 10 accepted for both human and animal use. During the entire immunisation period the total body weight of GDF-8 immunised as well as control animals will be monitored regularly. When the mice are approximately 16 weeks of age, they will be sacrificed and the size of their muscle mass will be determined.

15 Due to the relatively narrow window in time where mice are immune competent but still not fully outgrown, it may be difficult to demonstrate an effect of anti-GDF-8 vaccination in these animals. If that proves to be the case, it is contemplated to alternatively immunise rats or larger animals such as pigs, 20 cattle, etc.

The modified GDF-8 molecule, which superiorly increases the growth rate of the animals and/or superiorly increases the maximum muscle size, is selected for clinical development.

SEQUENCE LISTING

<110> M&E Biotech A/S

<120> Method for down-regulating GDF-8 activity

<130> AutoVacGDF-8 DK 1

<160> 23

<170> PatentIn Ver. 2.1

<210> 1

<211> 375

<212> PRT

<213> Homo sapiens

<400> 1

Met Gln Lys Leu Gln Leu Cys Val Tyr Ile Tyr Leu Phe Met Leu Ile
 1 5 10 15

Val Ala Gly Pro Val Asp Leu Asn Glu Asn Ser Glu Gln Lys Glu Asn
 20 25 30

Val Glu Lys Glu Gly Leu Cys Asn Ala Cys Thr Trp Arg Gln Asn Thr
 35 40 45

Lys Ser Ser Arg Ile Glu Ala Ile Lys Ile Gln Ile Leu Ser Lys Leu
 50 55 60

Arg Leu Glu Thr Ala Pro Asn Ile Ser Lys Asp Val Ile Arg Gln Leu
 65 70 75 80

Leu Pro Lys Ala Pro Pro Leu Arg Glu Leu Ile Asp Gln Tyr Asp Val
 85 90 95

Gln Arg Asp Asp Ser Ser Asp Gly Ser Leu Glu Asp Asp Asp Tyr His
 100 105 110

Ala Thr Thr Glu Thr Ile Ile Thr Met Pro Thr Glu Ser Asp Phe Leu
 115 120 125

Met Gln Val Asp Gly Lys Pro Lys Cys Cys Phe Phe Lys Phe Ser Ser
 130 135 140

Lys Ile Gln Tyr Asn Lys Val Val Lys Ala Gln Leu Trp Ile Tyr Leu
 145 150 155 160

Arg Pro Val Glu Thr Pro Thr Thr Val Phe Val Gln Ile Leu Arg Leu
 165 170 175

Ile Lys Pro Met Lys Asp Gly Thr Arg Tyr Thr Gly Ile Arg Ser Leu
 180 185 190

Lys Leu Asp Met Asn Pro Gly Thr Gly Ile Trp Gln Ser Ile Asp Val
 195 200 205
 Lys Thr Val Leu Gln Asn Trp Leu Lys Gln Pro Glu Ser Asn Leu Gly
 210 215 220
 Ile Glu Ile Lys Ala Leu Asp Glu Asn Gly His Asp Leu Ala Val Thr
 225 230 235 240
 Phe Pro Gly Pro Gly Glu Asp Gly Leu Asn Pro Phe Leu Glu Val Lys
 245 250 255
 Val Thr Asp Thr Pro Lys Arg Ser Arg Arg Asp Phe Gly Leu Asp Cys
 260 265 270
 Asp Glu His Ser Thr Glu Ser Arg Cys Cys Arg Tyr Pro Leu Thr Val
 275 280 285
 Asp Phe Glu Ala Phe Gly Trp Asp Trp Ile Ile Ala Pro Lys Arg Tyr
 290 295 300
 Lys Ala Asn Tyr Cys Ser Gly Glu Cys Glu Phe Val Phe Leu Gln Lys
 305 310 315 320
 Tyr Pro His Thr His Leu Val His Gln Ala Asn Pro Arg Gly Ser Ala
 325 330 335
 Gly Pro Cys Cys Thr Pro Thr Lys Met Ser Pro Ile Asn Met Leu Tyr
 340 345 350
 Phe Asn Gly Lys Glu Gln Ile Ile Tyr Gly Lys Ile Pro Ala Met Val
 355 360 365
 Val Asp Arg Cys Gly Cys Ser
 370 375

<210> 2
 <211> 362
 <212> PRT
 <213> Meleagris gallopavo

<400> 2
 Met Gln Ile Leu Val His Pro Val Ala Leu Asp Gly Ser Ser Gln Pro
 1 5 10 15

Thr Glu Asn Ala Glu Lys Asp Gly Leu Cys Asn Ala Cys Thr Trp Arg
 20 25 30
 Gln Asn Thr Lys Ser Ser Arg Ile Glu Ala Ile Lys Ile Gln Ile Leu
 35 40 45
 Ser Lys Leu Arg Leu Glu Gln Ala Pro Asn Ile Ser Arg Asp Val Ile
 50 55 60

Lys Gln Leu Leu Pro Lys Ala Pro Pro Leu Gln Glu Leu Ile Asp Gln
 65 70 75 80
 Tyr Asp Val Gln Arg Asp Asp Ser Ser Asp Gly Ser Leu Glu Asp Asp
 85 90 95
 Asp Tyr His Ala Thr Thr Glu Thr Ile Ile Thr Met Pro Thr Glu Ser
 100 105 110
 Asp Phe Leu Val Gln Met Glu Gly Lys Pro Lys Cys Cys Phe Phe Lys
 115 120 125
 Phe Ser Ser Lys Ile Gln Tyr Asn Lys Val Val Lys Ala Gln Leu Trp
 130 135 140
 Ile Tyr Leu Arg Gln Val Gln Lys Pro Thr Thr Val Phe Val Gln Ile
 145 150 155 160
 Leu Arg Leu Ile Lys Pro Met Lys Asp Gly Thr Arg Tyr Thr Gly Ile
 165 170 175
 Arg Ser Leu Lys Leu Asp Met Asn Pro Gly Thr Gly Ile Trp Gln Ser
 180 185 190
 Ile Asp Val Lys Thr Val Leu Gln Asn Trp Leu Lys Gln Pro Glu Ser
 195 200 205
 Asn Leu Gly Ile Glu Ile Lys Ala Phe Asp Glu Asn Gly Arg Asp Leu
 210 215 220
 Ala Val Thr Phe Pro Gly Pro Gly Glu Asp Gly Leu Asn Pro Phe Leu
 225 230 235 240
 Glu Val Arg Val Thr Asp Thr Pro Lys Arg Ser Arg Arg Asp Phe Gly
 245 250 255
 Leu Asp Cys Asp Glu His Ser Thr Glu Ser Arg Cys Cys Arg Tyr Pro
 260 265 270
 Leu Thr Val Asp Phe Glu Ala Phe Gly Trp Asp Trp Ile Ile Ala Pro
 275 280 285
 Lys Arg Tyr Lys Ala Asn Tyr Cys Ser Gly Glu Cys Glu Phe Val Phe
 290 295 300

 Leu Gln Lys Tyr Pro His Thr His Leu Val His Gln Ala Asn Pro Arg
 305 310 315 320
 Gly Ser Ala Gly Pro Cys Cys Thr Pro Thr Lys Met Ser Pro Ile Asn
 325 330 335
 Met Leu Tyr Phe Asn Gly Lys Glu Gln Ile Ile Tyr Gly Lys Ile Pro
 340 345 350

Ala Met Val Val Asp Arg Cys Gly Cys Ser
 355 360

<210> 3
 <211> 375
 <212> PRT
 <213> Gallus sp.

<400> 3
 Met Gln Lys Leu Ala Val Tyr Val Tyr Ile Tyr Leu Phe Met Gln Ile
 1 5 10 15
 Ala Val Asp Pro Val Ala Leu Asp Gly Ser Ser Gln Pro Thr Glu Asn
 20 25 30
 Ala Glu Lys Asp Gly Leu Cys Asn Ala Cys Thr Trp Arg Gln Asn Thr
 35 40 45
 Lys Ser Ser Arg Ile Glu Ala Ile Lys Ile Gln Ile Leu Ser Lys Leu
 50 55 60
 Arg Leu Glu Gln Ala Pro Asn Ile Ser Arg Asp Val Ile Lys Gln Leu
 65 70 75 80
 Leu Pro Lys Ala Pro Pro Leu Gln Glu Leu Ile Asp Gln Tyr Asp Val
 85 90 95
 Gln Arg Asp Asp Ser Ser Asp Gly Ser Leu Glu Asp Asp Asp Tyr His
 100 105 110
 Ala Thr Thr Glu Thr Ile Ile Thr Met Pro Thr Glu Ser Asp Phe Leu
 115 120 125
 Val Gln Met Glu Gly Lys Pro Lys Cys Cys Phe Phe Lys Phe Ser Ser
 130 135 140
 Lys Ile Gln Tyr Asn Lys Val Val Lys Ala Gln Leu Trp Ile Tyr Leu
 145 150 155 160
 Arg Gln Val Gln Lys Pro Thr Thr Val Phe Val Gln Ile Leu Arg Leu
 165 170 175
 Ile Lys Pro Met Lys Asp Gly Thr Arg Tyr Thr Gly Ile Arg Ser Leu
 180 185 190

Lys Leu Asp Met Asn Pro Gly Thr Gly Ile Trp Gln Ser Ile Asp Val
 195 200 205
 Lys Thr Val Leu Gln Asn Trp Leu Lys Gln Pro Glu Ser Asn Leu Gly
 210 215 220
 Ile Glu Ile Lys Ala Phe Asp Glu Thr Gly Arg Asp Leu Ala Val Thr
 225 230 235 240

Phe Pro Gly Pro Gly Glu Asp Gly Leu Asn Pro Phe Leu Glu Val Arg
245 250 255

Val Thr Asp Thr Pro Lys Arg Ser Arg Arg Asp Phe Gly Leu Asp Cys
260 265 270

Asp Glu His Ser Thr Glu Ser Arg Cys Cys Arg Tyr Pro Leu Thr Val
275 280 285

Asp Phe Glu Ala Phe Gly Trp Asp Trp Ile Ile Ala Pro Lys Arg Tyr
290 295 300

Lys Ala Asn Tyr Cys Ser Gly Glu Cys Glu Phe Val Phe Leu Gln Lys
305 310 315 320

Tyr Pro His Thr His Leu Val His Gln Ala Asn Pro Arg Gly Ser Ala
325 330 335

Gly Pro Cys Cys Thr Pro Thr Lys Met Ser Pro Ile Asn Met Leu Tyr
340 345 350

Phe Asn Gly Lys Glu Gln Ile Ile Tyr Gly Lys Ile Pro Ala Met Val
355 360 365

Val Asp Arg Cys Gly Cys Ser
370 375

<210> 4

<211> 376

<212> PRT

<213> Mus musculus

<400> 4

Met Met Gln Lys Leu Gln Met Tyr Val Tyr Ile Tyr Leu Phe Met Leu
1 5 10 15

Ile Ala Ala Gly Pro Val Asp Leu Asn Glu Gly Ser Glu Arg Glu Glu
20 25 30

Asn Val Glu Lys Glu Gly Leu Cys Asn Ala Cys Ala Trp Arg Gln Asn
35 40 45

Thr Arg Tyr Ser Arg Ile Glu Ala Ile Lys Ile Gln Ile Leu Ser Lys
50 55 60

Leu Arg Leu Glu Thr Ala Pro Asn Ile Ser Lys Asp Ala Ile Arg Gln
65 70 75 80

Leu Leu Pro Arg Ala Pro Pro Leu Arg Glu Leu Ile Asp Gln Tyr Asp
85 90 95

Val Gln Arg Asp Asp Ser Ser Asp Gly Ser Leu Glu Asp Asp Asp Tyr
100 105 110

His Ala Thr Thr Glu Thr Ile Ile Thr Met Pro Thr Glu Ser Asp Phe
 115 120 125
 Leu Met Gln Ala Asp Gly Lys Pro Lys Cys Cys Phe Phe Lys Phe Ser
 130 135 140
 Ser Lys Ile Gln Tyr Asn Lys Val Val Lys Ala Gln Leu Trp Ile Tyr
 145 150 155 160
 Leu Arg Pro Val Lys Thr Pro Thr Thr Val Phe Val Gln Ile Leu Arg
 165 170 175
 Leu Ile Lys Pro Met Lys Asp Gly Thr Arg Tyr Thr Gly Ile Arg Ser
 180 185 190
 Leu Lys Leu Asp Met Ser Pro Gly Thr Gly Ile Trp Gln Ser Ile Asp
 195 200 205
 Val Lys Thr Val Leu Gln Asn Trp Leu Lys Gln Pro Glu Ser Asn Leu
 210 215 220
 Gly Ile Glu Ile Lys Ala Leu Asp Glu Asn Gly His Asp Leu Ala Val
 225 230 235 240
 Thr Phe Pro Gly Pro Gly Glu Asp Gly Leu Asn Pro Phe Leu Glu Val
 245 250 255
 Lys Val Thr Asp Thr Pro Lys Arg Ser Arg Arg Asp Phe Gly Leu Asp
 260 265 270
 Cys Asp Glu His Ser Thr Glu Ser Arg Cys Cys Arg Tyr Pro Leu Thr
 275 280 285
 Val Asp Phe Glu Ala Phe Gly Trp Asp Trp Ile Ile Ala Pro Lys Arg
 290 295 300
 Tyr Lys Ala Asn Tyr Cys Ser Gly Glu Cys Glu Phe Val Phe Leu Gln
 305 310 315 320
 Lys Tyr Pro His Thr His Leu Val His Gln Ala Asn Pro Arg Gly Ser
 325 330 335
 Ala Gly Pro Cys Cys Thr Pro Thr Lys Met Ser Pro Ile Asn Met Leu
 340 345 350
 Tyr Phe Asn Gly Lys Glu Gln Ile Ile Tyr Gly Lys Ile Pro Ala Met
 355 360 365
 Val Val Asp Arg Cys Gly Cys Ser
 370 375

<210> 5
 <211> 375
 <212> PRT

<213> Bos taurus

<400> 5

Met Gln Lys Leu Gln Ile Ser Val Tyr Ile Tyr Leu Phe Met Leu Ile
 1 5 10 15

Val Ala Gly Pro Val Asp Leu Asn Glu Asn Ser Glu Gln Lys Glu Asn
 20 25 30

Val Glu Lys Glu Gly Leu Cys Asn Ala Cys Leu Trp Arg Glu Asn Thr
 35 40 45

Thr Ser Ser Arg Leu Glu Ala Ile Lys Ile Gln Ile Leu Ser Lys Leu
 50 55 60

Arg Leu Glu Thr Ala Pro Asn Ile Ser Lys Asp Ala Ile Arg Gln Leu
 65 70 75 80

Leu Pro Lys Ala Pro Pro Leu Leu Glu Leu Ile Asp Gln Phe Asp Val
 85 90 95

Gln Arg Asp Ala Ser Ser Asp Gly Ser Leu Glu Asp Asp Asp Tyr His
 100 105 110

Ala Arg Thr Glu Thr Val Ile Thr Met Pro Thr Glu Ser Asp Leu Leu
 115 120 125

Thr Gln Val Glu Gly Lys Pro Lys Cys Cys Phe Phe Lys Phe Ser Ser
 130 135 140

Lys Ile Gln Tyr Asn Lys Leu Val Lys Ala Gln Leu Trp Ile Tyr Leu
 145 150 155 160

Arg Pro Val Lys Thr Pro Ala Thr Val Phe Val Gln Ile Leu Arg Leu
 165 170 175

Ile Lys Pro Met Lys Asp Gly Thr Arg Tyr Thr Gly Ile Arg Ser Leu
 180 185 190

Lys Leu Asp Met Asn Pro Gly Thr Gly Ile Trp Gln Ser Ile Asp Val
 195 200 205

Lys Thr Val Leu Gln Asn Trp Leu Lys Gln Pro Glu Ser Asn Leu Gly
 210 215 220

Ile Glu Ile Lys Ala Leu Asp Glu Asn Gly His Asp Leu Ala Val Thr
 225 230 235 240

Phe Pro Glu Pro Gly Glu Asp Gly Leu Thr Pro Phe Leu Glu Val Lys
 245 250 255

Val Thr Asp Thr Pro Lys Arg Ser Arg Arg Asp Phe Gly Leu Asp Cys
 260 265 270

Asp Glu His Ser Thr Glu Ser Arg Cys Cys Arg Tyr Pro Leu Thr Val
 275 280 285
 Asp Phe Glu Ala Phe Gly Trp Asp Trp Ile Ile Ala Pro Lys Arg Tyr
 290 295 300
 Lys Ala Asn Tyr Cys Ser Gly Glu Cys Glu Phe Val Phe Leu Gln Lys
 305 310 315 320
 Tyr Pro His Thr His Leu Val His Gln Ala Asn Pro Arg Gly Ser Ala
 325 330 335
 Gly Pro Cys Cys Thr Pro Thr Lys Met Ser Pro Ile Asn Met Leu Tyr
 340 345 350
 Phe Asn Gly Glu Gly Gln Ile Ile Tyr Gly Lys Ile Pro Ala Met Val
 355 360 365
 Val Asp Arg Cys Gly Cys Ser
 370 375

<210> 6
 <211> 375
 <212> PRT
 <213> Ovis sp.

<400> 6
 Met Gln Lys Leu Gln Ile Phe Val Tyr Ile Tyr Leu Phe Met Leu Leu
 1 5 10 15
 Val Ala Gly Pro Val Asp Leu Asn Glu Asn Ser Glu Gln Lys Glu Asn
 20 25 30
 Val Glu Lys Lys Gly Leu Cys Asn Ala Cys Leu Trp Arg Gln Asn Asn
 35 40 45
 Lys Ser Ser Arg Leu Glu Ala Ile Lys Ile Gln Ile Leu Ser Lys Leu
 50 55 60
 Arg Leu Glu Thr Ala Pro Asn Ile Ser Lys Asp Ala Ile Arg Gln Leu
 65 70 75 80
 Leu Pro Lys Ala Pro Pro Leu Arg Glu Leu Ile Asp Gln Tyr Asp Val
 85 90 95

Gln Arg Asp Asp Ser Ser Asp Gly Ser Leu Glu Asp Asp Asp Tyr His
 100 105 110
 Val Thr Thr Glu Thr Val Ile Thr Met Pro Thr Glu Ser Asp Leu Leu
 115 120 125
 Ala Glu Val Gln Glu Lys Pro Lys Cys Cys Phe Phe Lys Phe Ser Ser
 130 135 140

Lys Ile Gln His Asn Lys Val Val Lys Ala Gln Leu Trp Ile Tyr Leu
 145 150 155 160
 Arg Pro Val Lys Thr Pro Thr Thr Val Phe Val Gln Ile Leu Arg Leu
 165 170 175
 Ile Lys Pro Met Lys Asp Gly Thr Arg Tyr Thr Gly Ile Arg Ser Leu
 180 185 190
 Lys Leu Asp Met Asn Pro Gly Thr Gly Ile Trp Gln Ser Ile Asp Val
 195 200 205
 Lys Thr Val Leu Gln Asn Trp Leu Lys Gln Pro Glu Ser Asn Leu Gly
 210 215 220
 Ile Glu Ile Lys Ala Leu Asp Glu Asn Gly His Asp Leu Ala Val Thr
 225 230 235 240
 Phe Pro Glu Pro Gly Glu Glu Gly Leu Asn Pro Phe Leu Glu Val Lys
 245 250 255
 Val Thr Asp Thr Pro Lys Arg Ser Arg Arg Asp Phe Gly Leu Asp Cys
 260 265 270
 Asp Glu His Ser Thr Glu Ser Arg Cys Cys Arg Tyr Pro Leu Thr Val
 275 280 285
 Asp Phe Glu Ala Phe Gly Trp Asp Trp Ile Ile Ala Pro Lys Arg Tyr
 290 295 300
 Lys Ala Asn Tyr Cys Ser Gly Glu Cys Glu Phe Leu Phe Leu Gln Lys
 305 310 315 320
 Tyr Pro His Thr His Leu Val His Gln Ala Asn Pro Lys Gly Ser Ala
 325 330 335
 Gly Pro Cys Cys Thr Pro Thr Lys Met Ser Pro Ile Asn Met Leu Tyr
 340 345 350
 Phe Asn Gly Lys Glu Gln Ile Ile Tyr Gly Lys Ile Pro Gly Met Val
 355 360 365
 Val Asp Arg Cys Gly Cys Ser
 370 375

<210> 7
 <211> 376
 <212> PRT
 <213> Rattus norvegicus

<400> 7
 Met Ile Gln Lys Pro Gln Met Tyr Val Tyr Ile Tyr Leu Phe Val Leu
 1 5 10 15

Ile	Ala	Ala	Gly	Pro	Val	Asp	Leu	Asn	Glu	Asp	Ser	Glu	Arg	Glu	Ala	20	25	30
Asn	Val	Glu	Lys	Glu	Gly	Leu	Cys	Asn	Ala	Cys	Ala	Trp	Arg	Gln	Asn	35	40	45
Thr	Arg	Tyr	Ser	Arg	Ile	Glu	Ala	Ile	Lys	Ile	Gln	Ile	Leu	Ser	Lys	50	55	60
Leu	Arg	Leu	Glu	Thr	Ala	Pro	Asn	Ile	Ser	Lys	Asp	Ala	Ile	Arg	Gln	65	70	75
Leu	Leu	Pro	Arg	Ala	Pro	Pro	Leu	Arg	Glu	Leu	Ile	Asp	Gln	Tyr	Asp	85	90	95
Val	Gln	Arg	Asp	Asp	Ser	Ser	Asp	Gly	Ser	Leu	Glu	Asp	Asp	Asp	Tyr	100	105	110
His	Ala	Thr	Thr	Glu	Thr	Ile	Ile	Thr	Met	Pro	Thr	Glu	Ser	Asp	Phe	115	120	125
Leu	Met	Gln	Ala	Asp	Gly	Lys	Pro	Lys	Cys	Cys	Phe	Phe	Lys	Phe	Ser	130	135	140
Ser	Lys	Ile	Gln	Tyr	Asn	Lys	Val	Val	Lys	Ala	Gln	Leu	Trp	Ile	Tyr	145	150	155
Leu	Arg	Ala	Val	Lys	Thr	Pro	Thr	Thr	Val	Phe	Val	Gln	Ile	Leu	Arg	165	170	175
Leu	Ile	Lys	Pro	Met	Lys	Asp	Gly	Thr	Arg	Tyr	Thr	Gly	Ile	Arg	Ser	180	185	190
Leu	Lys	Leu	Asp	Met	Ser	Pro	Gly	Thr	Gly	Ile	Trp	Gln	Ser	Ile	Asp	195	200	205
Val	Lys	Thr	Val	Leu	Gln	Asn	Trp	Leu	Lys	Gln	Pro	Glu	Ser	Asn	Leu	210	215	220
Gly	Ile	Glu	Ile	Lys	Ala	Leu	Asp	Glu	Asn	Gly	His	Asp	Leu	Ala	Val	225	230	235
Thr	Phe	Pro	Gly	Pro	Gly	Glu	Asp	Gly	Leu	Asn	Pro	Phe	Leu	Glu	Val	245	250	255
Lys	Val	Thr	Asp	Thr	Pro	Lys	Arg	Ser	Arg	Arg	Asp	Phe	Gly	Leu	Asp	260	265	270
Cys	Asp	Glu	His	Ser	Thr	Glu	Ser	Arg	Cys	Cys	Arg	Tyr	Pro	Leu	Thr	275	280	285
Val	Asp	Phe	Glu	Ala	Phe	Gly	Trp	Asp	Trp	Ile	Ile	Ala	Pro	Lys	Arg	290	295	300

Tyr Lys Ala Asn Tyr Cys Ser Gly Glu Cys Glu Phe Val Phe Leu Gln
 305 310 315 320

Lys Tyr Pro His Thr His Leu Val His Gln Ala Asn Pro Arg Gly Ser
 325 330 335

Ala Gly Pro Cys Cys Thr Pro Thr Lys Met Ser Pro Ile Asn Met Leu
 340 345 350

Tyr Phe Asn Gly Lys Glu Gln Ile Ile Tyr Gly Lys Ile Pro Ala Met
 355 360 365

Val Val Asp Arg Cys Gly Cys Ser
 370 375

<210> 8

<211> 375

<212> PRT

<213> Sus scrofa

<400> 8

Met Gln Lys Leu Gln Ile Tyr Val Tyr Ile Tyr Leu Phe Met Leu Ile
 1 5 10 15

Val Ala Gly Pro Val Asp Leu Asn Glu Asn Ser Glu Gln Lys Glu Asn
 20 25 30

Val Glu Lys Glu Gly Leu Cys Asn Ala Cys Met Trp Arg Gln Asn Thr
 35 40 45

Lys Ser Ser Arg Leu Glu Ala Ile Lys Ile Gln Ile Leu Ser Lys Leu
 50 55 60

Arg Leu Glu Thr Ala Pro Asn Ile Ser Lys Asp Ala Ile Arg Gln Leu
 65 70 75 80

Leu Pro Lys Ala Pro Pro Leu Arg Glu Leu Ile Asp Gln Tyr Asp Val
 85 90 95

Gln Arg Asp Asp Ser Ser Asp Gly Ser Leu Glu Asp Asp Asp Tyr His
 100 105 110

Ala Thr Thr Glu Thr Ile Ile Thr Met Pro Thr Glu Ser Asp Leu Leu
 115 120 125

Met Gln Val Glu Gly Lys Pro Lys Cys Cys Phe Phe Lys Phe Ser Ser
 130 135 140

Lys Ile Gln Tyr Asn Lys Val Val Lys Ala Gln Leu Trp Ile Tyr Leu
 145 150 155 160

Arg Pro Val Lys Thr Pro Thr Thr Val Phe Val Gln Ile Leu Arg Leu
 165 170 175

Ile Lys Pro Met Lys Asp Gly Thr Arg Tyr Thr Gly Ile Arg Ser Leu
 180 185 190
 Lys Leu Asp Met Asn Pro Gly Thr Gly Ile Trp Gln Ser Ile Asp Val
 195 200 205
 Lys Thr Val Leu Gln Asn Trp Leu Lys Gln Pro Glu Ser Asn Leu Gly
 210 215 220
 Ile Glu Ile Lys Ala Leu Asp Glu Asn Gly His Asp Leu Ala Val Thr
 225 230 235 240
 Phe Pro Gly Pro Gly Glu Asp Gly Leu Asn Pro Phe Leu Glu Val Lys
 245 250 255
 Val Thr Asp Thr Pro Lys Arg Ser Arg Arg Asp Phe Gly Leu Asp Cys
 260 265 270
 Asp Glu His Ser Thr Glu Ser Arg Cys Cys Arg Tyr Pro Leu Thr Val
 275 280 285
 Asp Phe Glu Ala Phe Gly Trp Asp Trp Ile Ile Ala Pro Lys Arg Tyr
 290 295 300
 Lys Ala Asn Tyr Cys Ser Gly Glu Cys Glu Phe Val Phe Leu Gln Lys
 305 310 315 320
 Tyr Pro His Thr His Leu Val His Gln Ala Asn Pro Arg Gly Ser Ala
 325 330 335
 Gly Pro Cys Cys Thr Pro Thr Lys Met Ser Pro Ile Asn Met Leu Tyr
 340 345 350
 Phe Asn Gly Lys Glu Gln Ile Ile Tyr Gly Lys Ile Pro Ala Met Val
 355 360 365
 Val Asp Arg Cys Gly Cys Ser
 370 375

<210> 9
 <211> 374
 <212> PRT
 <213> Danio rerio

<400> 9

Met His Phe Thr Gln Val Leu Ile Ser Leu Ser Val Leu Ile Ala Cys
 1 5 10 15
 Gly Pro Val Gly Tyr Gly Asp Ile Thr Ala His Gln Gln Pro Ser Thr
 20 25 30
 Ala Thr Glu Glu Ser Glu Leu Cys Ser Thr Cys Glu Phe Arg Gln His
 35 40 45

Ser Lys Leu Met Arg Leu His Ala Ile Lys Ser Gln Ile Leu Ser Lys
 50 55 60
 Leu Arg Leu Lys Gln Ala Pro Asn Ile Ser Arg Asp Val Val Lys Gln
 65 70 75 80
 Leu Leu Pro Lys Ala Pro Pro Leu Gln Gln Leu Leu Asp Gln Tyr Asp
 85 90 95
 Val Leu Gly Asp Asp Ser Lys Asp Gly Ala Val Glu Glu Asp Asp Glu
 100 105 110
 His Ala Thr Thr Glu Thr Ile Met Thr Met Ala Thr Glu Pro Asp Pro
 115 120 125
 Ile Val Gln Val Asp Arg Lys Pro Lys Cys Cys Phe Phe Ser Phe Ser
 130 135 140
 Pro Lys Ile Gln Ala Asn Arg Ile Val Arg Ala Gln Leu Trp Val His
 145 150 155 160
 Leu Arg Pro Ala Glu Glu Ala Thr Thr Val Phe Leu Gln Ile Ser Arg
 165 170 175
 Leu Met Pro Val Lys Asp Gly Gly Arg His Arg Ile Arg Ser Leu Lys
 180 185 190
 Ile Asp Val Asn Ala Gly Val Thr Ser Trp Gln Ser Ile Asp Val Lys
 195 200 205
 Gln Val Leu Thr Val Trp Leu Lys Gln Pro Glu Thr Asn Arg Gly Ile
 210 215 220
 Glu Ile Asn Ala Tyr Asp Ala Lys Gly Asn Asp Leu Ala Val Thr Ser
 225 230 235 240
 Thr Glu Thr Gly Glu Asp Gly Leu Leu Pro Phe Met Glu Val Lys Ile
 245 250 255
 Ser Glu Gly Pro Lys Arg Ile Arg Arg Asp Ser Gly Leu Asp Cys Asp
 260 265 270
 Glu Asn Ser Ser Glu Ser Arg Cys Cys Arg Tyr Pro Leu Thr Val Asp
 275 280 285

 Phe Glu Asp Phe Gly Trp Asp Trp Ile Ile Ala Pro Lys Arg Tyr Lys
 290 295 300
 Ala Asn Tyr Cys Ser Gly Glu Cys Asp Tyr Met Tyr Leu Gln Lys Tyr
 305 310 315 320
 Pro His Thr His Leu Val Asn Lys Ala Ser Pro Arg Gly Thr Ala Gly
 325 330 335

Pro Cys Cys Thr Pro Thr Lys Met Ser Pro Ile Asn Met Leu Tyr Phe
 340 345 350

Asn Gly Lys Glu Gln Ile Ile Tyr Gly Lys Ile Pro Ser Met Val Val
 355 360 365

Asp Arg Cys Gly Cys Ser
 370

<210> 10

<211> 375

<212> PRT

<213> Papio hamadryas

<400> 10

Met Gln Lys Leu Gln Leu Cys Val Tyr Ile Tyr Leu Phe Met Leu Ile
 1 5 10 15

Val Ala Gly Pro Val Asp Leu Asn Glu Asn Ser Glu Gln Lys Glu Asn
 20 25 30

Val Glu Lys Glu Gly Leu Cys Asn Ala Cys Thr Trp Arg Gln Asn Thr
 35 40 45

Lys Ser Ser Arg Ile Glu Ala Ile Lys Ile Gln Ile Leu Ser Lys Leu
 50 55 60

Arg Leu Glu Thr Ala Pro Asn Ile Ser Lys Asp Ala Ile Arg Gln Leu
 65 70 75 80

Leu Pro Lys Ala Pro Pro Leu Arg Glu Leu Ile Asp Gln Tyr Asp Val
 85 90 95

Gln Arg Asp Asp Ser Ser Asp Gly Ser Leu Glu Asp Asp Asp Tyr His
 100 105 110

Ala Thr Thr Glu Thr Ile Ile Thr Met Pro Thr Glu Ser Asp Phe Leu
 115 120 125

Met Gln Val Asp Gly Lys Pro Lys Cys Cys Phe Phe Lys Phe Ser Ser
 130 135 140

Lys Ile Gln Tyr Asn Lys Val Val Lys Ala Gln Leu Trp Ile Tyr Leu
 145 150 155 160

Arg Pro Val Glu Thr Pro Thr Thr Val Phe Val Gln Ile Leu Arg Leu
 165 170 175

Ile Lys Pro Met Lys Asp Gly Thr Arg Tyr Thr Gly Ile Arg Ser Leu
 180 185 190

Lys Leu Asp Met Asn Pro Gly Thr Gly Ile Trp Gln Ser Ile Asp Val
 195 200 205

Lys Thr Val Leu Gln Asn Trp Leu Lys Gln Pro Glu Ser Asn Leu Gly
 210 215 220

Ile Glu Ile Lys Ala Leu Asp Glu Asn Gly His Asp Leu Ala Val Thr
 225 230 235 240

Phe Pro Gly Pro Gly Glu Asp Gly Leu Asn Pro Phe Leu Glu Val Lys
 245 250 255

Val Thr Asp Thr Pro Lys Arg Ser Arg Arg Asp Phe Gly Leu Asp Cys
 260 265 270

Asp Glu His Ser Thr Glu Ser Arg Cys Cys Arg Tyr Pro Leu Thr Val
 275 280 285

Asp Phe Glu Ala Leu Gly Trp Asp Trp Ile Ile Ala Pro Lys Arg Tyr
 290 295 300

Lys Ala Asn Tyr Cys Ser Gly Glu Cys Glu Phe Val Phe Leu Gln Lys
 305 310 315 320

Tyr Pro His Thr His Leu Val His Gln Ala Asn Pro Arg Gly Ser Ala
 325 330 335

Gly Pro Cys Cys Thr Pro Thr Lys Met Ser Pro Ile Asn Met Leu Tyr
 340 345 350

Phe Asn Gly Lys Glu Gln Ile Ile Tyr Gly Lys Ile Pro Ala Met Val
 355 360 365

Val Asp Arg Cys Gly Cys Ser
 370 375

<210> 11

<211> 109

<212> PRT

<213> Homo sapiens

<220>

<221> PEPTIDE

<222> (1)..(109)

<223> Identical to residues 267-375 in SEQ ID NO: 1

<400> 11

Asp Phe Gly Leu Asp Cys Asp Glu His Ser Thr Glu Ser Arg Cys Cys
 1 5 10 15

Arg Tyr Pro Leu Thr Val Asp Phe Glu Ala Phe Gly Trp Asp Trp Ile
 20 25 30

Ile Ala Pro Lys Arg Tyr Lys Ala Asn Tyr Cys Ser Gly Glu Cys Glu
 35 40 45

Phe Val Phe Leu Gln Lys Tyr Pro His Thr His Leu Val His Gln Ala
 50 55 60

Asn Pro Arg Gly Ser Ala Gly Pro Cys Cys Thr Pro Thr Lys Met Ser
 65 70 75 80

Pro Ile Asn Met Leu Tyr Phe Asn Gly Lys Glu Gln Ile Ile Tyr Gly
 85 90 95

Lys Ile Pro Ala Met Val Val Asp Arg Cys Gly Cys Ser
 100 105

<210> 12
 <211> 109
 <212> PRT
 <213> Bos taurus

<220>
 <221> PEPTIDE
 <222> (1)..(109)
 <223> Identical to residues 267-375 in SEQ ID NO: 5

<400> 12
 Asp Phe Gly Leu Asp Cys Asp Glu His Ser Thr Glu Ser Arg Cys Cys
 1 5 10 15

Arg Tyr Pro Leu Thr Val Asp Phe Glu Ala Phe Gly Trp Asp Trp Ile
 20 25 30

Ile Ala Pro Lys Arg Tyr Lys Ala Asn Tyr Cys Ser Gly Glu Cys Glu
 35 40 45

Phe Val Phe Leu Gln Lys Tyr Pro His Thr His Leu Val His Gln Ala
 50 55 60

Asn Pro Arg Gly Ser Ala Gly Pro Cys Cys Thr Pro Thr Lys Met Ser
 65 70 75 80

Pro Ile Asn Met Leu Tyr Phe Asn Gly Glu Gly Gln Ile Ile Tyr Gly
 85 90 95

Lys Ile Pro Ala Met Val Val Asp Arg Cys Gly Cys Ser
 100 105

<210> 13
 <211> 15
 <212> PRT
 <213> Clostridium tetani

<400> 13
 Gln Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly Ile Thr Glu Leu
 1 5 10 15

<210> 14
 <211> 21
 <212> PRT
 <213> Clostridium tetani

<400> 14
 Phe Asn Asn Phe Thr Val Ser Phe Trp Leu Arg Val Pro Lys Val Ser
 1 5 10 15
 Ala Ser His Leu Glu
 20

<210> 15
 <211> 109
 <212> PRT
 <213> Artificial sequence

<220>
 <221> MUTAGEN
 <222> (18)..(32)
 <223> Tetanus toxoid P2 epitope (SEQ ID NO: 13)

<220>
 <221> SIMILAR
 <222> (1)..(17)
 <223> Identical to residues 267-283 in SEQ ID NO: 1

<220>
 <221> SIMILAR
 <222> (33)..(109)
 <223> Identical to residues 299-375 in SEQ ID NO: 1

<220>
 <221> SITE
 <222> (73)
 <223> Cys or Ser

<220>
 <221> SITE
 <222> (90)..(91)
 <223> Lys Glu or Glu Gly

<400> 15
 Asp Phe Gly Leu Asp Cys Asp Glu His Ser Thr Glu Ser Arg Cys Cys
 1 5 10 15

Arg Gln Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly Ile Thr Glu Leu
 20 25 30

Ile Ala Pro Lys Arg Tyr Lys Ala Asn Tyr Cys Ser Gly Glu Cys Glu
 35 40 45

Phe Val Phe Leu Gln Lys Tyr Pro His Thr His Leu Val His Gln Ala
 50 55 60

Asn Pro Arg Gly Ser Ala Gly Pro Cys Cys Thr Pro Thr Lys Met Ser
 65 70 75 80

Pro Ile Asn Met Leu Tyr Phe Asn Gly Lys Glu Gln Ile Ile Tyr Gly
 85 90 95

Lys Ile Pro Ala Met Val Val Asp Arg Cys Gly Cys Ser
 100 105

<210> 16

<211> 109

<212> PRT

<213> Artificial sequence

<220>

<221> MUTAGEN

<222> (52)..(66)

<223> Tetanus toxoid P2 epitope (SEQ ID NO: 13)

<220>

<221> SIMILAR

<222> (1)..(51)

<223> Identical to residues 267-317 in SEQ ID NO: 1

<220>

<221> SIMILAR

<222> (67)..(109)

<223> Identical to residues 333-375 in SEQ ID NO: 1

<220>

<221> SITE

<222> (73)

<223> Cys or Ser

<220>

<221> SITE

<222> (90)..(91)

<223> Lys Glu or Glu Gly

<400> 16

Asp Phe Gly Leu Asp Cys Asp Glu His Ser Thr Glu Ser Arg Cys Cys
 1 5 10 15

Arg Tyr Pro Leu Thr Val Asp Phe Glu Ala Phe Gly Trp Asp Trp Ile
 20 25 30

Ile Ala Pro Lys Arg Tyr Lys Ala Asn Tyr Cys Ser Gly Glu Cys Glu
 35 40 45

Phe Val Phe Gln Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly Ile Thr
 50 55 60

Glu Leu Arg Gly Ser Ala Gly Pro Cys Cys Thr Pro Thr Lys Met Ser
 65 70 75 80

Pro Ile Asn Met Leu Tyr Phe Asn Gly Lys Glu Gln Ile Ile Tyr Gly
 85 90 95

Lys Ile Pro Ala Met Val Val Asp Arg Cys Gly Cys Ser
 100 105

<210> 17
 <211> 109
 <212> PRT
 <213> Artificial sequence

<220>
 <221> MUTAGEN
 <222> (83)..(97)
 <223> Tetanus toxoid P2 epitope (SEQ ID NO: 13)

<220>
 <221> SIMILAR
 <222> (1)..(82)
 <223> Identical to residues 267-348 in SEQ ID NO: 1

<220>
 <221> SIMILAR
 <222> (98)..(109)
 <223> Identical to residues 364-375 in SEQ ID NO: 1

<220>
 <221> SITE
 <222> (73)
 <223> Cys or Ser

<220>
 <221> SITE
 <222> (90)..(91)
 <223> Lys Glu or Glu Gly

<400> 17
 Asp Phe Gly Leu Asp Cys Asp Glu His Ser Thr Glu Ser Arg Cys Cys
 1 5 10 15

Arg Tyr Pro Leu Thr Val Asp Phe Glu Ala Phe Gly Trp Asp Trp Ile
 20 25 30

Ile Ala Pro Lys Arg Tyr Lys Ala Asn Tyr Cys Ser Gly Glu Cys Glu
 35 40 45

Phe Val Phe Leu Gln Lys Tyr Pro His Thr His Leu Val His Gln Ala
 50 55 60

Asn Pro Arg Gly Ser Ala Gly Pro Cys Cys Thr Pro Thr Lys Met Ser
 65 70 75 80

Pro Ile Gln Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly Ile Thr Glu
 85 90 95

Leu Ile Pro Ala Met Val Val Asp Arg Cys Gly Cys Ser
 100 105

<210> 18
 <211> 109
 <212> PRT
 <213> Artificial sequence

<220>
 <221> MUTAGEN
 <222> (21)..(41)
 <223> Tetanus toxoid P30 epitope (SEQ ID NO: 14)

<220>
 <221> SIMILAR
 <222> (42)..(109)
 <223> Identical to residues 307-375 in SEQ ID NO: 1

<220>
 <221> SIMILAR
 <222> (42)..(109)
 <223> Identical to residues 308-375 in SEQ ID NO: 1

<220>
 <221> SITE
 <222> (73)
 <223> Cys or Ser

<220>
 <221> SITE
 <222> (90)..(91)
 <223> Lys Glu or Glu Gly

<400> 18
 Asp Phe Gly Leu Asp Cys Asp Glu His Ser Thr Glu Ser Arg Cys Cys
 1 5 10 15

Arg Tyr Pro Leu Phe Asn Asn Phe Thr Val Ser Phe Trp Leu Arg Val
 20 25 30

Pro Lys Val Ser Ala Ser His Leu Glu Tyr Cys Ser Gly Glu Cys Glu
 35 40 45

Phe Val Phe Leu Gln Lys Tyr Pro His Thr His Leu Val His Gln Ala
 50 55 60

Asn Pro Arg Gly Ser Ala Gly Pro Cys Cys Thr Pro Thr Lys Met Ser
 65 70 75 80

Pro Ile Asn Met Leu Tyr Phe Asn Gly Lys Glu Gln Ile Ile Tyr Gly
 85 90 95

Lys Ile Pro Ala Met Val Val Asp Arg Cys Gly Cys Ser
 100 105

<210> 19
 <211> 109
 <212> PRT
 <213> Artificial sequence

<220>
 <221> MUTAGEN
 <222> (49)..(69)
 <223> Tetanus toxoid P30 epitope (SEQ ID NO: 14)

<220>
 <221> SIMILAR
 <222> (1)..(48)
 <223> Identical to residues 267-314 in SEQ ID NO: 1

<220>
 <221> SIMILAR
 <222> (70)..(109)
 <223> Identical to residues 336-375 in SEQ ID NO: 1

<220>
 <221> SITE
 <222> (73)
 <223> Cys or Ser

<220>
 <221> SITE
 <222> (90)..(91)
 <223> Lys Glu or Glu Gly

<400> 19
 Asp Phe Gly Leu Asp Cys Asp Glu His Ser Thr Glu Ser Arg Cys Cys
 1 5 10 15
 Arg Tyr Pro Leu Thr Val Asp Phe Glu Ala Phe Gly Trp Asp Trp Ile
 20 25 30
 Ile Ala Pro Lys Arg Tyr Lys Ala Asn Tyr Cys Ser Gly Glu Cys Glu
 35 40 45
 Phe Asn Asn Phe Thr Val Ser Phe Trp Leu Arg Val Pro Lys Val Ser
 50 55 60
 Ala Ser His Leu Glu Ala Gly Pro Cys Cys Thr Pro Thr Lys Met Ser
 65 70 75 80
 Pro Ile Asn Met Leu Tyr Phe Asn Gly Lys Glu Gln Ile Ile Tyr Gly
 85 90 95
 Lys Ile Pro Ala Met Val Val Asp Arg Cys Gly Cys Ser
 100 105

<210> 20

<211> 109
 <212> PRT
 <213> Artificial sequence

<220>
 <221> MUTAGEN
 <222> (79)..(99)
 <223> Tetanus toxoid P30 epitope (SEQ ID NO: 14)

<220>
 <221> SIMILAR
 <222> (1)..(78)
 <223> Identical to residues 267-345 in SEQ ID NO: 1

<220>
 <221> SIMILAR
 <222> (100)..(109)
 <223> Identical to residues 366-375 in SEQ ID NO: 1

<220>
 <221> SITE
 <222> (73)
 <223> Cys or Ser

<220>
 <221> SITE
 <222> (90)..(91)
 <223> Lys Glu or Glu Gly

<400> 20
 Asp Phe Gly Leu Asp Cys Asp Glu His Ser Thr Glu Ser Arg Cys Cys
 1 5 10 15

Arg Tyr Pro Leu Thr Val Asp Phe Glu Ala Phe Gly Trp Asp Trp Ile
 20 25 30

Ile Ala Pro Lys Arg Tyr Lys Ala Asn Tyr Cys Ser Gly Glu Cys Glu
 35 40 45

Phe Val Phe Leu Gln Lys Tyr Pro His Thr His Leu Val His Gln Ala
 50 55 60

Asn Pro Arg Gly Ser Ala Gly Pro Cys Cys Thr Pro Thr Lys Phe Asn
 65 70 75 80

Asn Phe Thr Val Ser Phe Trp Leu Arg Val Pro Lys Val Ser Ala Ser
 85 90 95

His Leu Glu Ala Met Val Val Asp Arg Cys Gly Cys Ser
 100 105

<210> 21
 <211> 109
 <212> PRT

<213> Artificial sequence

<220>

<221> MUTAGEN

<222> (84)..(104)

<223> Tetanus toxoid P30 epitope (SEQ ID NO: 14)

<220>

<221> SIMILAR

<222> (1)..(83)

<223> Identical to residues 267-349 in SEQ ID NO: 1

<220>

<221> SIMILAR

<222> (105)..(109)

<223> Identical to residues 371-375 in SEQ ID NO: 1

<220>

<221> SITE

<222> (73)

<223> Cys or Ser

<220>

<221> SITE

<222> (90)..(91)

<223> Lys Glu or Glu Gly

<400> 21

Asp	Phe	Gly	Leu	Asp	Cys	Asp	Glu	His	Ser	Thr	Glu	Ser	Arg	Cys	Cys
1				5					10					15	

Arg	Tyr	Pro	Leu	Thr	Val	Asp	Phe	Glu	Ala	Phe	Gly	Trp	Asp	Trp	Ile
			20					25					30		

Ile	Ala	Pro	Lys	Arg	Tyr	Lys	Ala	Asn	Tyr	Cys	Ser	Gly	Glu	Cys	Glu
	35						40					45			

Phe	Val	Phe	Leu	Gln	Lys	Tyr	Pro	His	Thr	His	Leu	Val	His	Gln	Ala
	50				55						60				

Asn	Pro	Arg	Gly	Ser	Ala	Gly	Pro	Cys	Cys	Thr	Pro	Thr	Lys	Met	Ser
65					70					75				80	

Pro	Ile	Asn	Phe	Asn	Asn	Phe	Thr	Val	Ser	Phe	Trp	Leu	Arg	Val	Pro
			85						90					95	

Lys	Val	Ser	Ala	Ser	His	Leu	Glu	Arg	Cys	Gly	Cys	Ser
			100					105				

<210> 22

<211> 254

<212> PRT

<213> Artificial sequence

<220>
 <221> SIMILAR
 <222> (110)..(124)
 <223> Tetanus toxoid P2 epitope (SEQ ID NO: 13)

<220>
 <221> SIMILAR
 <222> (125)..(145)
 <223> Diphtheria toxoid P30 epitope (SEQ ID NO: 14)

<220>
 <221> SIMILAR
 <222> (1)..(109)
 <223> 109 C-terminal residues of human and bovine GDF-8
 (residues 267-375 in SEQ ID NO: 1)

<220>
 <221> SIMILAR
 <222> (146)..(254)
 <223> 109 C-terminal residues of human and bovine GDF-8
 (residues 267-375 in SEQ ID NO: 1)

<220>
 <221> SITE
 <222> (90)..(91)
 <223> Lys Glu or Glu Gly

<220>
 <221> SITE
 <222> (235)..(236)
 <223> Identical to (90)..(91)

<400> 22
 Asp Phe Gly Leu Asp Cys Asp Glu His Ser Thr Glu Ser Arg Cys Cys
 1 5 10 15
 Arg Tyr Pro Leu Thr Val Asp Phe Glu Ala Phe Gly Trp Asp Trp Ile
 20 25 30
 Ile Ala Pro Lys Arg Tyr Lys Ala Asn Tyr Cys Ser Gly Glu Cys Glu
 35 40 45
 Phe Val Phe Leu Gln Lys Tyr Pro His Thr His Leu Val His Gln Ala
 50 55 60

Asn Pro Arg Gly Ser Ala Gly Pro Cys Cys Thr Pro Thr Lys Met Ser
 65 70 75 80
 Pro Ile Asn Met Leu Tyr Phe Asn Gly Lys Glu Gln Ile Ile Tyr Gly
 85 90 95
 Lys Ile Pro Ala Met Val Val Asp Arg Cys Gly Cys Ser Gln Tyr Ile
 100 105 110

Lys Ala Asn Ser Lys Phe Ile Gly Ile Thr Glu Leu Phe Asn Asn Phe
 115 120 125
 Thr Val Ser Phe Trp Leu Arg Val Pro Lys Val Ser Ala Ser His Leu
 130 135 140
 Glu Asp Phe Gly Leu Asp Cys Asp Glu His Ser Thr Glu Ser Arg Cys
 145 150 155 160
 Cys Arg Tyr Pro Leu Thr Val Asp Phe Glu Ala Phe Gly Trp Asp Trp
 165 170 175
 Ile Ile Ala Pro Lys Arg Tyr Lys Ala Asn Tyr Cys Ser Gly Glu Cys
 180 185 190
 Glu Phe Val Phe Leu Gln Lys Tyr Pro His Thr His Leu Val His Gln
 195 200 205
 Ala Asn Pro Arg Gly Ser Ala Gly Pro Cys Cys Thr Pro Thr Lys Met
 210 215 220
 Ser Pro Ile Asn Met Leu Tyr Phe Asn Gly Lys Glu Gln Ile Ile Tyr
 225 230 235 240
 Gly Lys Ile Pro Ala Met Val Val Asp Arg Cys Gly Cys Ser
 245 250

<210> 23
 <211> 160
 <212> PRT
 <213> Artificial sequence

<220>
 <221> MUTAGEN
 <222> (16)..(36)
 <223> Tetanus toxoid P30 epitope (SEQ ID NO: 14)

<220>
 <221> MUTAGEN
 <222> (37)..(51)
 <223> Tetanus toxoid P2 epitope (SEQ ID NO: 13)

<220>
 <221> SIMILAR
 <222> (1)..(15)
 <223> Identical to residues 216-230 of SEQ ID NO: 1

<220>
 <221> SIMILAR
 <222> (52)..(160)
 <223> Identical to residues 267-375 of SEQ ID NO: 1

<220>
 <221> SITE

<222> (124)

<223> Cys or Ser

<220>

<221> SITE

<222> (141)..(142)

<223> Lys Glu or Glu Gly

<400> 23

Leu	Lys	Gln	Pro	Glu	Ser	Asn	Leu	Gly	Ile	Glu	Ile	Lys	Ala	Leu	Phe
1				5				10						15	

Asn	Asn	Phe	Thr	Val	Ser	Phe	Trp	Leu	Arg	Val	Pro	Lys	Val	Ser	Ala
			20					25					30		

Ser	His	Leu	Glu	Gln	Tyr	Ile	Lys	Ala	Asn	Ser	Lys	Phe	Ile	Gly	Ile
		35					40					45			

Thr	Glu	Leu	Asp	Phe	Gly	Leu	Asp	Cys	Asp	Glu	His	Ser	Thr	Glu	Ser
	50					55					60				

Arg	Cys	Cys	Arg	Tyr	Pro	Leu	Thr	Val	Asp	Phe	Glu	Ala	Phe	Gly	Trp
65					70					75					80

Asp	Trp	Ile	Ile	Ala	Pro	Lys	Arg	Tyr	Lys	Ala	Asn	Tyr	Cys	Ser	Gly
				85					90					95	

Glu	Cys	Glu	Phe	Val	Phe	Leu	Gln	Lys	Tyr	Pro	His	Thr	His	Leu	Val
			100					105					110		

His	Gln	Ala	Asn	Pro	Arg	Gly	Ser	Ala	Gly	Pro	Cys	Cys	Thr	Pro	Thr
		115					120					125			

Lys	Met	Ser	Pro	Ile	Asn	Met	Leu	Tyr	Phe	Asn	Gly	Lys	Glu	Gln	Ile
	130					135					140				

Ile	Tyr	Gly	Lys	Ile	Pro	Ala	Met	Val	Val	Asp	Arg	Cys	Gly	Cys	Ser
145					150					155					160

CLAIMS

1. A method for *in vivo* down-regulation of growth differentiation factor 8 (GDF-8) activity in an animal, including a human being, the method comprising effecting presentation to the animal's immune system of an immunogenically effective amount of
 - at least one GDF-8 polypeptide or subsequence thereof which has been formulated so that immunization of the animal with the GDF-8 polypeptide or subsequence thereof induces production of antibodies against the GDF-8 polypeptide, and/or
 - at least one GDF-8 analogue wherein is introduced at least one modification in the GDF-8 amino acid sequence which has as a result that immunization of the animal with the analogue induces production of antibodies against the GDF-8 polypeptide.
2. The method according to claim 1, wherein is presented a GDF-8 analogue with at least one modification of the GDF-8 amino acid sequence.
3. The method according to claim 2, wherein the modification has as a result that a substantial fraction of GDF-8 B-cell epitopes are preserved and that
 - at least one foreign T helper lymphocyte epitope (T_H epitope) is introduced, and/or
 - at least one first moiety is introduced which effects targeting of the modified molecule to an antigen presenting cell (APC) or a B-lymphocyte, and/or
 - at least one second moiety is introduced which stimulates the immune system, and/or
 - at least one third moiety is introduced which optimizes presentation of the modified GDF-8 polypeptide to the immune system.

4. The method according to claim 3, wherein the modification includes introduction as side groups, by covalent or non-covalent binding to suitable chemical groups in GDF-8 or a subsequence thereof, of the foreign T_H epitope and/or of the first and/or of the second and/or of the third moiety.
5. The method according to claim 3 or 4, wherein the modification includes amino acid substitution and/or deletion and/or insertion and/or addition.
6. The method according to claim 5, wherein the modification results in the provision of a fusion polypeptide.
7. The method according to claim 5 or 6, wherein introduction of the amino acid substitution and/or deletion and/or insertion and/or addition results in a substantial preservation of the overall tertiary structure of GDF-8.
8. The method according to any one of claims 2-7, wherein the modification includes duplication of at least one GDF-8 B-cell epitope and/or introduction of a hapten.
9. The method according to any one of claims 3-8, wherein the foreign T-cell epitope is immunodominant in the animal.
10. The method according to any one of claims 3-9, wherein the foreign T-cell epitope is promiscuous.
11. The method according to claim 10, wherein the at least one foreign T-cell epitope is selected from a natural promiscuous T-cell epitope and an artificial MHC-II binding peptide sequence.
12. The method according to claim 11, wherein the natural T-cell epitope is selected from a Tetanus toxoid epitope such as P2 or P30, a diphtheria toxoid epitope, an influenza virus hemagglutinin epitope, and a *P. falciparum* CS epitope.

13. The method according to any one of claims 3-12, wherein the first moiety is a substantially specific binding partner for a B-lymphocyte specific surface antigen or for an APC specific surface antigen.

5 14. The method according to claim 13, wherein the first moiety is a carbohydrate for which there is a receptor on the B-lymphocyte or the APC, such as mannan or mannose or wherein the first moiety is a hapten.

15. The method according to any one of claims 3-14, wherein the
10 second moiety is selected from a cytokine, a hormone, and a heat-shock protein.

16. The method according to claim 6, wherein the cytokine is selected from, or is an effective part of, interferon γ (IFN- γ), Flt3L, interleukin 1 (IL-1), interleukin 2 (IL-2), interleukin 4
15 (IL-4), interleukin 6 (IL-6), interleukin 12 (IL-12), interleukin 13 (IL-13), interleukin 15 (IL-15), and granulocyte-macrophage colony stimulating factor (GM-CSF).

17. The method according to claim 15, wherein the heat-shock protein is selected from the group consisting of HSP70, HSP90,
20 HSC70, GRP94, and calreticulin (CRT), or an effective part thereof.

18. The method according to any one of claims 3-17, wherein the third moiety is of lipid nature, such as a palmitoyl group, a myristyl group, a farnesyl group, a geranyl-geranyl group, a
25 GPI-anchor, and an N-acyl diglyceride group.

19. The method according to any one of the preceding claims, wherein the GDF-8 subsequence or the GDF-8 analogue is derived from the C-terminal, active form of GDF-8.

20. The method according to claim 19, wherein the GDF-8 subsequence or GDF-8 analogue is derived from a bovine, porcine, human, chicken, sheep, or turkey GDF-8 polypeptide.

21. The method according to claim 20, wherein the GDF-8 polypeptide has been modified by substituting at least one amino acid sequence SEQ ID NO: 11 or 12 with at least one amino acid sequence of equal or different length and containing a foreign T_H epitope, wherein substituted amino acid sequences are comprised in residues 18-41, 49-69, or 79-104 in SEQ ID NO: 11 or 12.

22. The method according to any one of the preceding claims, wherein presentation to the immune system is effected by having at least two copies of the GDF-8 polypeptide, the subsequence thereof or the modified GDF-8 polypeptide covalently or non-covalently linked to a carrier molecule capable of effecting presentation of multiple copies of antigenic determinants.

23. The method according to any the preceding claims, wherein the GDF-8 polypeptide, the subsequence thereof, or the modified GDF-8 polypeptide has been formulated with an adjuvant which facilitates breaking of autotolerance to autoantigens.

24. The method according to claim 23, wherein the adjuvant is selected from the group consisting of an immune targeting adjuvant; an immune modulating adjuvant such as a toxin, a cytokine and a mycobacterial derivative; an oil formulation; a polymer; a micelle forming adjuvant; a saponin; an immunostimulating complex matrix (an ISCOM matrix); a particle; DDA; aluminium adjuvants; DNA adjuvants; γ -inulin; and an encapsulating adjuvant.

25. The method according to any one of the preceding claims, wherein an effective amount of the GDF-8 polypeptide or the GDF-8 analogue is administered to the animal via a route

selected from the parenteral route such as the intradermal, the subdermal, the intracutaneous, the subcutaneous, and the intramuscular routes; the peritoneal route; the oral route; the buccal route; the sublingual route; the epidural route; the spinal route; the anal route; and the intracranial route.

26. The method according to claim 25, wherein the effective amount is between 0.5 μ g and 2,000 μ g of the GDF-8 polypeptide, the subsequence thereof or the analogue thereof.

27. The method according to claim 25 or 26, which includes at least one administration of the GDF-8 polypeptide or analogue per year, such as at least 2, at least 3, at least 4, at least 6, and at least 12 administrations per year.

28. The method according to any one of claims 25-27, wherein the GDF-8 polypeptide or analogue is contained in a virtual lymph node (VLN) device.

29. The method according to any one of claims 1-24, wherein presentation of modified GDF-8 to the immune system is effected by introducing nucleic acid(s) encoding the modified GDF-8 into the animal's cells and thereby obtaining *in vivo* expression by the cells of the nucleic acid(s) introduced.

30. The method according to claim 29, wherein the nucleic acid(s) introduced is/are selected from naked DNA, DNA formulated with charged or uncharged lipids, DNA formulated in liposomes, DNA included in a viral vector, DNA formulated with a transfection-facilitating protein or polypeptide, DNA formulated with a targeting protein or polypeptide, DNA formulated with Calcium precipitating agents, DNA coupled to an inert carrier molecule, DNA formulated with chitin or chitosan, and DNA formulated with an adjuvant.

31. The method according to claim 29 or 30, wherein the nucleic acids are administered intraarterially, intravenously, or by the routes defined in claim 25.

32. The method according to claim 30 or 31, wherein the nucleic acid(s) is/are contained in a VLN device.

33. The method according to any one of claims 30-32, which includes at least one administration of the nucleic acids per year, such as at least 2, at least 3, at least 4, at least 6, and at least 12 administrations per year

34. A method for increasing the muscle mass of an animal, the method comprising down-regulating GDF-8 activity according to the method of any one of claims 1-33 to such an extent such that the muscle mass is increased at least 5% when compared to animals which exhibit normal GDF-8 activity, such as at least 10, 15, 20, 25, 30, 35, 40, and 45%.

35. A GDF-8 analogue which is derived from an animal GDF-8 polypeptide wherein is introduced a modification which has as a result that immunization of the animal with the analogue induces production of antibodies against the GDF-8 polypeptide.

36. A GDF-8 analogue according to claim 35, wherein the modification is as defined in any one of claims 1-22.

37. An immunogenic composition comprising an immunogenically effective amount of a GDF-8 polypeptide autologous in an animal, said GDF-8 polypeptide being formulated together with an

immunologically acceptable adjuvant so as to break the animal's autotolerance towards the GDF-8 polypeptide, the composition further comprising a pharmaceutically and immunologically acceptable carrier and/or vehicle.

38. An immunogenic composition comprising an immunogenically effective amount of an GDF-8 analogue according to claim 35 or 36, the composition further comprising a pharmaceutically and immunologically acceptable carrier and/or vehicle and optionally
5 an adjuvant.

39. An immunogenic composition according to Claim 37 or 38, wherein the adjuvant is selected from the group consisting of the adjuvants of claim 24.

40. A nucleic acid fragment which encodes an GDF-8 analogue
10 according to claim 35 or 36.

41. A vector carrying the nucleic acid fragment according to claim 40.

42. The vector according to claim 41 which is capable of autonomous replication.

15 43. The vector according to claim 41 or 42 which is selected from the group consisting of a plasmid, a phage, a cosmid, a mini-chromosome, and a virus.

44. The vector according to any one of claims 41-43, comprising, in the 5'-3' direction and in operable linkage, a promoter for
20 driving expression of the nucleic acid fragment according to claim 40, optionally a nucleic acid sequence encoding a leader peptide enabling secretion of or integration into the membrane of the polypeptide fragment, the nucleic acid fragment according to claim 40, and optionally a terminator.

25 45. The vector according to any one of claims 41-44 which, when introduced into a host cell, is integrated in the host cell genome.

46. The vector according to any one of claims 41-44 which, when introduced into a host cell, is not capable of being integrated in the host cell genome.

47. The vector according to any one of claims 41-46, wherein the promoter drives expression in a eukaryotic cell.

48. The vector according to any one of claims 41-46, wherein the promoter drives expression in a prokaryotic cell.

49. A transformed cell carrying the vector of any one of claims 41-48.

50. The transformed cell according to claim 49 which is capable of replicating the nucleic acid fragment according to claim 40.

51. The transformed cell according to claim 50, which is a microorganism selected from a bacterium, a yeast, a protozoan, or a cell derived from a multicellular organism selected from a fungus, an insect cell such as an S₂ or an SF cell, a plant cell, and a mammalian cell.

52. The transformed cell according to claim 51 which is a bacterium of the genus *Escherichia*, *Bacillus*, *Salmonella*, or *Mycobacterium*.

53. The transformed cell according to claim 52, which is an *E. coli* cell.

54. The transformed cell according to claim 53, which is a non-pathogenic *Mycobacterium* cell such as *M. bovis* BCG.

55. The transformed cell according to any one of claims 49-54, which expresses the nucleic acid fragment according to claim 40.

56. The transformed cell according to claim 55, which secretes or carries on its surface, the GDF-8 analogue according to claim 35 or 36.

57. The method according to any one of claims 1-22, wherein presentation to the immune system is effected by administering a non-pathogenic microorganism or virus which is carrying a nucleic acid fragment which encodes and expresses the GDF-8 polypeptide or analogue.

58. The method according to claim 57, wherein the virus is a non-virulent pox virus such as a vaccinia virus.

59. The method according to claim 58, wherein the microorganism is a bacterium.

60. The method according to claim 58, wherein the bacterium is as defined in any one of claims 52-54.

61. The method according to any one of claims 57-60, wherein the non-pathogenic microorganism or virus is administered one single time to the animal.

62. A composition for inducing production of antibodies against GDF-8, the composition comprising

- a nucleic acid fragment according to claim 40 or a vector according to any one of claims 41-48, and
- a pharmaceutically and immunologically acceptable carrier and/or vehicle and/or adjuvant.

63. The composition according to claim 62, wherein the nucleic acid fragment is formulated according to claim 30 or 32.

64. A stable cell line which carries the vector according to any one of claims 41-48 and which expresses the nucleic acid fragment according to claim 40, and which optionally secretes or

carries the GDF-8 analogue according to claim 35 or 36 on its surface.

65. A method for the preparation of the cell according to any one of claims 49-56, the method comprising transforming a host cell with the nucleic acid fragment according to claim 40 or with the vector according to any one of claims 41-48.

66. A method for the identification of a modified GDF-8 polypeptide which is capable of inducing antibodies against unmodified GDF-8 in an animal species where the unmodified GDF-8 polypeptide is a self-protein, the method comprising

- preparing, by means of peptide synthesis or genetic engineering techniques, a set of mutually distinct modified GDF-8 polypeptides wherein amino acids have been added to, inserted in, deleted from, or substituted into the amino acid sequence of an GDF-8 polypeptide of the animal species thereby giving rise to amino acid sequences in the set which comprise T-cell epitopes which are foreign to the animal species,
- testing members of the set for their ability to induce production of antibodies by the animal species against the unmodified GDF-8, and
- isolating the member(s) of the set which significantly induces antibody production against unmodified GDF-8 in the animal species.

67. A method for the preparation of an immunogenic composition comprising at least one modified GDF-8 polypeptide which is capable of inducing antibodies against unmodified GDF-8 in an animal species where the unmodified GDF-8 polypeptide is a self-protein, the method comprising

- preparing, by means of peptide synthesis or genetic engineering techniques, a set of mutually distinct modified

GDF-8 polypeptides wherein amino acids have been added to, inserted in, deleted from, or substituted into the amino acid sequence of an GDF-8 polypeptide of the animal species thereby giving rise to amino acid sequences in the set comprising T-cell epitopes which are foreign to the animal, testing members of the set for their ability to induce production of antibodies by the animal species against the unmodified GDF-8, and and admixing the member(s) of the set which significantly induces production of antibodies in the animal species which are reactive with GDF-8 with a pharmaceutically and immunologically acceptable carrier and/or vehicle, optionally in combination with at least one pharmaceutically and immunologically acceptable adjuvant.

68. The method according to claim 66 or 67, wherein preparation of the members of the set comprises preparation of mutually distinct nucleic acid sequences, each sequence being a nucleic acid sequence according to claim 40, insertion of the nucleic acid sequences into appropriate expression vectors, transformation of suitable host cells with the vectors, and expression of the nucleic acid sequences, optionally followed by isolation of the expression products.

69. The method according to claim 68, wherein the preparation of the nucleic acid sequences and/or the vectors is achieved by the aid of a molecular amplification technique such as PCR

70. The method according to claim 68, wherein the preparation of the nucleic sequences and/or the vectors is achieved by the aid of nucleic acid synthesis.

71. Use of GDF-8 or a subsequence thereof for the preparation of an immunogenic composition comprising an adjuvant for down-regulating GDF-8 activity in an animal.

72. Use of GDF-8 or a subsequence thereof for the preparation of an immunogenic composition comprising an adjuvant for increasing the muscle mass of an animal.

73. Use of an GDF-8 analogue for the preparation of an
5 immunogenic composition optionally comprising an adjuvant for down-regulating GDF-8 activity in an animal.

74. Use of an GDF-8 analogue for the preparation of an immunogenic composition optionally comprising an adjuvant for increasing the muscle mass of an animal.



Fig. 1A



Fig. 1B



Fig. 1C




Fig. 1D




Fig. 2

